

Asymptotic Relative Entropy of Entanglement

K. Audenaert,¹ J. Eisert,^{2,3} E. Jané,^{2,4} M.B. Plenio,² S. Virmani,² and B. De Moor¹

¹Department of Electrical Engineering (ESAT-SISTA), Katholieke Universiteit Leuven, B-3001 Leuven-Heverlee, Belgium

²QOLS, Blackett Laboratory, Imperial College of Science, Technology and Medicine, London SW7 2BW, United Kingdom

³Institut für Physik, Universität Potsdam, 14469 Potsdam, Germany

⁴Departament d'Estructura i Constituents de la Matèria, Universitat de Barcelona, E-08028 Barcelona, Spain

(Received 17 April 2001; published 6 November 2001)

We present an analytical formula for the asymptotic relative entropy of entanglement with respect to positive partial transpose states for Werner states of arbitrary dimension. We then demonstrate its validity using methods from convex optimization. This is the first case in which the asymptotic value of a subadditive entanglement measure has been calculated.

DOI: 10.1103/PhysRevLett.87.217902

PACS numbers: 03.65.Ud, 03.67.Hk

The systematic investigation of quantum entanglement is a major goal of quantum information theory [1]. In the study of entanglement manipulation one considers the interconversion of different forms of entanglement by means of local quantum operations and classical communication (LQCC). For pure bipartite states entanglement manipulation in both the finite and asymptotic limits is quite well understood. For pure states necessary and sufficient conditions for the local interconvertibility of entangled states are known. In the asymptotic limit of infinitely many copies of a pure state, a single number, the von Neumann entropy of a subsystem, appropriately quantifies the degree of entanglement [2].

Much less is known about the entanglement of mixed states. One approach is to define *entanglement measures*, which are functions of a state that cannot increase under local operations and provide constraints on possible local entanglement manipulation protocols. These measures prove to be useful mathematical and conceptual tools and have interesting links with other areas such as the study of channel capacities [3]. A number of such measures have been proposed, most notably the entanglement of formation [3], the distillable entanglement [3,4], and the relative entropy of entanglement [4,5]. The distillable entanglement is defined as the asymptotic number of pure maximally entangled states that can be obtained via LQCC from a supply of a given state. For mixed states the distillable entanglement is exceedingly difficult to compute as it is defined as an asymptotic quantity referring to infinitely many copies of a quantum state. Therefore, upper bounds on the distillable entanglement, in particular, other entanglement measures, are of major practical use. One such entanglement measure is the relative entropy of entanglement, defined as

$$E_R(\sigma) = \min_{\rho \in \mathcal{D}} S(\sigma || \rho), \quad (1)$$

for states σ , where \mathcal{D} is the set of states with positive partial transpose (PPT states), and $S(\sigma || \rho) = \text{tr}[\sigma \lg \sigma - \sigma \lg \rho]$ (\lg signifies \log_2). This function essentially quantifies the distinguishability of σ from the set of PPT states. The set \mathcal{D} can also be taken to be the set of separable states [5]. However, the set of PPT states is much easier to

characterize, and the resulting measure provides a tighter bound to the distillable entanglement, one that is actually attained on pure states and certain mixed states [3].

In general, efficient protocols for entanglement manipulation require an asymptotic number of copies of the initial state. Therefore, to address any question related to asymptotic entanglement manipulation, one will instead have to consider asymptotic versions of the entanglement measures. For a given measure of entanglement E , the asymptotic version E^∞ is defined as the average entanglement per copy in the asymptotic limit [6,7],

$$E^\infty(\sigma) = \lim_{n \rightarrow \infty} \frac{E(\sigma^{\otimes n})}{n}. \quad (2)$$

For example, the asymptotic cost of creating a mixed state by LQCC from a supply of pure maximally entangled states is given by the asymptotic entanglement of formation [7]. However, such asymptotic entanglement measures are difficult to compute, and so far this task has not been accomplished except for the very rare occasions where the measure in question is known to be additive [8].

In this Letter we present a general formula for the asymptotic relative entropy of entanglement for an important class of bipartite states, namely, the Werner states of arbitrary dimension [9]. These states are the only




FIG. 1. The asymptotic relative entropy of entanglement $E_R^\infty(\sigma(p))$ of Werner states in $\mathbb{C}^d \otimes \mathbb{C}^d$ as a function of p (the weight of the antisymmetric state) for several values of d . The dashed lines separate the two regimes of Eq. (3), for $d = 10$ and for $d = 3$.

states that are invariant under local unitaries of the form $U \otimes U$. They can be conveniently formulated in terms of the projectors π_0 (π_1) on the symmetric (antisymmetric) subspaces of a Hilbert space $\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d$. Denoting the permutation operator that interchanges the subsystems as π , the projectors can be expressed as $\pi_0 = (1 + \pi)/2$ and $\pi_1 = (1 - \pi)/2$. A general Werner state is of the form $\sigma(p) = p\sigma_1 + (1 - p)\sigma_0$, $p \in [0, 1]$, where $\sigma_0 = \pi_0/\text{tr}[\pi_0]$ and $\sigma_1 = \pi_1/\text{tr}[\pi_1]$. Appropriately exploiting the symmetry of this state is one of the key ingredients to the proof of the main statement of this Letter. We formulate this main result as a theorem.

Theorem: For states in $\mathbb{C}^d \otimes \mathbb{C}^d$ of the form $\sigma(p) = p\sigma_1 + (1 - p)\sigma_0$ with $p \in (1/2, 1]$, E_R^∞ with respect to (w.r.t.) PPT states is given by

$$E_R^\infty[\sigma(p)] = \begin{cases} 1 - H(p), & p \leq \frac{d+2}{2d} \\ \lg \frac{d+2}{d} + (1-p) \lg \frac{d-2}{d+2}, & p > \frac{d+2}{2d} \end{cases} \quad (3)$$

where $H(p) = -p \lg p - (1-p) \lg(1-p)$.

It is interesting to note from this formula that E_R^∞ never exceeds the logarithm of the negativity [10]. Curiously, $E_R^\infty(\sigma(p))$ is a convex function of p , whereas numerics show that $E_R(\sigma(p)^{\otimes n})/n$ is not convex for any finite number $n > 1$. It is also intriguing that this formula is *exactly* the same as that from a different optimization problem considered by Rains [11], where he minimizes the function $B(\rho, \sigma) = S(\rho|\sigma) + \log|\sigma^\Gamma|$ over *all* states σ . Rains's quantity is also an upper bound to the distillable entanglement. However, it is not a convex function, which makes its minimization difficult due to the possibility of local minima. This applies especially in the asymptotic limit as the additivity properties of $B(\rho, \sigma)$ are unknown.

Proof: The proof proceeds in two stages. First we provide an upper bound on $E_R(\sigma^{\otimes n})/n$ for any number of copies n , by presenting a trial optimal state. Then we derive a lower bound for $E_R(\sigma^{\otimes n})/n$ using convex optimization methods. We conclude by showing that the two bounds coincide in the asymptotic limit $n \rightarrow \infty$.

Upper bound.—We consider a situation where we hold n copies of a Werner state $\sigma(p)$. Following the work of Ref. [12], we make heavy use of symmetry. In Refs. [4,12] it has been shown that if a state is invariant under a certain symmetry group, then one can restrict the minimization in Eq. (1) to those PPT states that are also invariant under the same group. As $\sigma(p)$ is invariant under the group G of local unitaries of the form $U \otimes U$, the state $\sigma(p)^{\otimes n}$ is invariant under $G^{\otimes n}$. This implies that the optimal PPT state for $\sigma(p)^{\otimes n}$ can be chosen to be a convex sum of the 2^n possible n -fold tensor products of σ_0 and σ_1 ,

$$\eta = \sum_{f=0}^{2^n-1} x_f (\sigma_{f_1} \otimes \sigma_{f_2} \otimes \dots \otimes \sigma_{f_n}), \quad (4)$$

where $\vec{x} = (x_0, \dots, x_{2^n-1})^T$ forms a probability distribution, and $f_i \in \{0, 1\}$, $i = 1, \dots, n$, is the i th digit in the binary representation of f . Furthermore, as the state $\sigma^{\otimes n}$ is invariant under any permutation of the n copies, we can

add the constraint that $x_l = x_m$ if the number of 1's in the binary representations of l and m are the same.

We consider trial states of the form $\sum_i w_i (u_i \sigma_1 + v_i \sigma_0)^{\otimes n}$, which automatically satisfy this constraint. Such states correspond to

$$\vec{x} = \sum_i w_i \begin{pmatrix} u_i \\ v_i \end{pmatrix}^{\otimes n}, \quad (5)$$

where the $u_i(v_i)$ component is the weight of $\sigma_1(\sigma_0)$, $u_i + v_i = 1$, and $\sum_i w_i = 1$ (and therefore $\sum_j x_j = 1$).

We need to know the eigenvalues of the partial transpose of η in order to ensure that it is PPT. Ignoring degeneracy, there are only two eigenvalues of the partial transpose of the Werner state $\sigma(p)$. It is easy to show that they are non-negative if and only if (iff) the following two-component vector is non-negative:

$$\begin{pmatrix} -1 & 1 \\ 1 & (d-1)/(d+1) \end{pmatrix} \begin{pmatrix} p \\ 1-p \end{pmatrix} = T \begin{pmatrix} p \\ 1-p \end{pmatrix}. \quad (6)$$

Similarly, it can easily be shown that η will be a PPT state iff the \vec{y} and \vec{x} in the following equation are non-negative vectors:

$$\vec{y} = \begin{pmatrix} -1 & 1 \\ 1 & (d-1)/(d+1) \end{pmatrix}^{\otimes n} \vec{x} = T^{\otimes n} \vec{x}. \quad (7)$$

Subject to this condition we need to calculate the relative entropy between the n -copy Werner state $\sigma(p)^{\otimes n}$ and this generalized Werner state η . This is given by

$$\begin{aligned} S(\sigma(p)^{\otimes n} || \eta) / n &= -H(p) \\ &\quad - (1/n) \sum_{k=0}^n C_n^k p^{n-k} (1-p)^k \\ &\quad \times \lg \sum_i w_i u_i^{n-k} v_i^k. \end{aligned} \quad (8)$$

At this point, we notice that since the second term in Eq. (8) is the average of the function

$$\xi(n-k) = (1/n) \lg \sum_i w_i u_i^{n-k} v_i^k \quad (9)$$

over a binomially distributed variable k , the value can be substituted by $\xi(np)$ when we take the limit $n \rightarrow \infty$ [13].

Since $E_R^\infty(\sigma(p))$ is the minimal value of $S(\sigma(p)^{\otimes n} || \eta) / n$ over all possible PPT states η , any such PPT state η will give us an upper bound for $E_R^\infty(\sigma(p))$. In particular, for the vector \vec{x} we propose a mixture of two n -fold Kronecker powers:

$$\vec{x} = \sum_{i=1}^2 w_i \begin{pmatrix} 1 - a_i \\ a_i \end{pmatrix}^{\otimes n}, \quad (10)$$

where $0 \leq w_i \leq 1$, $w_1 = 1 - w_2$, $0 \leq a_1 \leq 1/2$, and $1/2 \leq a_2 \leq 1$. Proper values for the parameters need to be selected to ensure that the corresponding state η will be PPT. Inspired by numerical results, we consider two separate intervals for p in Eq. (3): $1/2 \leq p \leq (d+2)/(2d)$ and $(d+2)/(2d) \leq p \leq 1$.

For the first interval, set $w_1 = 0$ and $a_2 = 1/2$ so that $x_k = 2^{-n}$ for all k . This state gives us an upper bound that equals E_R for one copy of $\sigma(p)$:

$$E_R^\infty(\sigma(p)) \leq \frac{S(\sigma(p)^{\otimes n} \parallel \eta)}{n} = 1 - H(p). \quad (11)$$

Now consider the second interval. We will set

$$a_1 = \frac{(d+2)(1-p)}{d+2-4p}, \quad a_2 = \frac{1+d-(d+2)a_1}{d+2-4a_1}, \quad (12)$$

$$w_1 = 1 - w_2 = \frac{1}{1+z^n}, \quad z = (d+2-4a_1)/d.$$

We calculate $T^{\otimes n} \vec{x}$:

$$\vec{y} = T^{\otimes n} \vec{x} = \sum_{i=1}^2 w_i \left(\begin{array}{c} 2a_i - 1 \\ 1 - 2a_i/(d+1) \end{array} \right)^{\otimes n} \quad (13)$$

so that, with the values of Eq. (12)

$$y_k = \sum_{i=1}^2 w_i (2a_i - 1)^{n-\#(k)} \left(1 - \frac{2a_i}{d+1} \right)^{\#(k)}, \quad (14)$$

where $\#(k)$ is the number of 1's in the binary representation of k . It is easy to check using Eq. (12) that y_k is always non-negative.

As a consequence, by taking into account the discussion after Eq. (8) and the notation introduced in Eq. (5), the upper bound for $n \rightarrow \infty$ reads

$$\begin{aligned} E_R^\infty(\sigma(p)) &\leq -H(p) - \lim_{n \rightarrow \infty} \lg \left(\sum_{i=1}^2 w_i u_i^{pn} v_i^{(1-p)n} \right)^{1/n} \\ &= -H(p) - \lg \lim_{n \rightarrow \infty} \left(\frac{t_1^n + t_2^n}{q_1^n + q_2^n} \right)^{1/n} \\ &= -H(p) - \lg \frac{\max(t_1, t_2)}{\max(q_1, q_2)} \end{aligned} \quad (15)$$

with

$$\begin{aligned} t_1 &= d(d-2)^p (d+2)^{1-p} (1-p)^{1-p} p^p, \\ t_2 &= [d-2+d^2(1-p)]^p (d^2 p - d - 2)^{1-p}, \\ q_1 &= d^2 - 4 \quad \text{and} \quad q_2 = d(d+2-4p). \end{aligned} \quad (16)$$

It is easy to check that for $p \geq (d+2)/(2d)$ (the second interval), both $t_1 \geq t_2$ and $q_1 \geq q_2$. So we obtain

$$\begin{aligned} E_R^\infty(\sigma(p)) &\leq -H(p) - \lg \frac{t_1}{q_1} \\ &= \lg \frac{d-2}{d} + p \lg \frac{d+2}{d-2}. \end{aligned} \quad (17)$$

In the other regime, $p \leq (d+2)/(2d)$, the bound obtainable from Eq. (15) is worse than Eq. (11). This ends the proof of the upper bound.

Proof of lower bound.—We now proceed to find a lower bound on E_R . To do this, we use the idea of Lagrange duality [14]. To calculate E_R^∞ we need to solve the optimization problem

$$\begin{aligned} \frac{E_R(\sigma(p)^{\otimes n})}{n} &= \min_{\vec{x}} \left\{ -H(p) - \frac{1}{n} \vec{z}^T \cdot \lg \vec{x} \right\} \\ \text{with } \vec{x} &\geq 0, \quad -T^{\otimes n} \vec{x} \leq 0, \quad \sum_{k=0}^{2^n-1} x_k = 1, \end{aligned} \quad (18)$$

where $\vec{z}^T = (p, 1-p)^{\otimes n}$. This is a convex optimization problem, so it is possible to consider the so-called dual problem. It is a basic result in convex optimization theory that the optimal (maximal) g value of the dual problem is a lower bound on the optimal (minimal) value of the primal problem, which is just what we are looking for (see [14] for a general description of duality in optimization).

The dual problem can be obtained as follows. First form the Lagrangian by multiplying the constraints with Lagrange multipliers and adding them to the objective function:

$$\begin{aligned} \mathcal{L}(\vec{x}, \vec{\lambda}, \nu) &= -H(p) - \frac{1}{n} \vec{z}^T \lg \vec{x} \\ &\quad - \vec{\lambda}^T T^{\otimes n} \vec{x} + \nu \left(\sum_{k=0}^{2^n-1} x_k - 1 \right). \end{aligned} \quad (19)$$

The constraint $\vec{x} \geq 0$ is not included explicitly, as it determines just the domain of the function \lg . Note that $\vec{\lambda}$ must exhibit the same copy-permutation symmetry as \vec{x} . Because the constraint associated to $\vec{\lambda}$ is an inequality, we have to introduce the constraint $\vec{\lambda} \geq 0$. The dual function is now given by

$$g(\vec{\lambda}, \nu) = \inf_{\vec{x} \geq 0} \mathcal{L}(\vec{x}, \vec{\lambda}, \nu), \quad (20)$$

and the dual optimization problem is

$$\max_{\vec{\lambda}, \nu} g(\vec{\lambda}, \nu), \quad \vec{\lambda} \geq 0, \quad (21)$$

including any other constraints on the domain of g .

For our problem, the dual function can be calculated explicitly. The derivative of the Lagrangian w.r.t. x_k is

$$\frac{\partial \mathcal{L}}{\partial x_k} = -\frac{1}{n \ln 2} \frac{z_k}{x_k} - (\vec{\lambda}^T \cdot T^{\otimes n})_k + \nu. \quad (22)$$

The Lagrangian reaches an extremum (minimum) at

$$\hat{x}_k = \frac{1}{\mu_k} \frac{z_k}{n \ln 2}, \quad (23)$$

where $\mu_k = \nu - (T^{\otimes n} \vec{\lambda})_k$ and we have exploited the symmetry of T . The dual function is

$$g(\vec{\lambda}, \nu) = \frac{1 + \ln(n \ln 2)}{n \ln 2} + \frac{1}{n} \sum_{k=0}^{2^n-1} z_k \lg \mu_k - \nu, \quad (24)$$

where we have used that $\vec{z}^T = (p, 1-p)^{\otimes n}$, which implies that $\sum_{k=0}^{2^n-1} z_k \lg z_k = -nH(p)$. As stated before, $\vec{\lambda}$ must be non-negative, and inspecting the domain of $g(\vec{\lambda}, \nu)$ yields an additional constraint that the μ_k be non-negative.

Now, any acceptable assignment of values to $\vec{\lambda}$ and ν gives a lower bound to $E_R^\infty(\sigma(p))$. Again we consider the two p intervals of Eq. (3). For $p \leq (d+2)/(2d)$ we propose

$$\begin{aligned} \nu^1 &= 1/(n \ln 2), \\ \vec{\lambda}^1 &= \frac{\nu^1}{d^n} \left[\begin{pmatrix} 1 \\ d+1 \end{pmatrix}^{\otimes n} - \begin{pmatrix} d+1-2dp \\ d+1 \end{pmatrix}^{\otimes n} \right]. \end{aligned} \quad (25)$$

After a short calculation we obtain $\mu_k = \nu^1 2^n p^{n-\#(k)} (1-p)^{\#(k)}$. This gives a feasible point of the dual problem, because both $\tilde{\lambda}$ and μ_k as given here are non-negative. The value of g using these assignments is $g(\tilde{\lambda}^1, \nu^1) = 1 - H(p)$.

For the second interval, $p \geq p' = (d+2)/(2d)$, we replace p by p' in Eq. (25), giving

$$\nu^2 = \nu^1, \tilde{\lambda}^2 = \frac{\nu^2}{d^n} \left[\begin{pmatrix} 1 \\ d+1 \end{pmatrix}^{\otimes n} - \begin{pmatrix} -1 \\ d+1 \end{pmatrix}^{\otimes n} \right]. \quad (26)$$

We now obtain $g(\tilde{\lambda}^2, \nu^2) = \lg \frac{d+2}{d} + (1-p) \lg \frac{d-2}{d+2}$.

As the two lower bounds $g(\tilde{\lambda}^1, \nu^1)$ and $g(\tilde{\lambda}^2, \nu^2)$ coincide with the two upper bounds [Eqs. (11) and (17)], the proof of the Theorem is now complete. \square

The remarkable behavior of $E_R^\infty(\sigma(p))$ is shown in Fig. 1 for several values of d . The nonlinear behavior for small values of p goes over into a linear dependence on p above the threshold value $p' = (d+2)/2d$. An immediate consequence of the result is that there are no inseparable Werner states with zero entanglement cost—a similar conclusion could not be drawn from Rains's bound as it is not an asymptotic quantity. It is astonishing that, as long as $p \leq p'$, E_R^∞ is invariant under the strongly irreversible operation of twirling, mapping Werner states on $\mathbb{C}^2 \otimes \mathbb{C}^2$ to Werner states on $\mathbb{C}^d \otimes \mathbb{C}^d$ [15].

Interestingly, the dependence of $E_R^\infty(\sigma(p))$ on p is quite similar to the conjectured behavior of the entanglement of formation for a single copy of an isotropic state [16]: there, one can also distinguish between two regimes, and for larger values of the weight F of the maximally entangled state in the isotropic state the dependence of the entanglement of formation is conjectured to be linearly dependent on F .

In this Letter we have concentrated on the important class of Werner states. With similar methods, one can also investigate other classes with high symmetry. It is hoped that this work can significantly contribute to the quest for a better understanding of the asymptotic regime of entanglement.

We thank R. F. Werner and D. Jonathan for fruitful discussions and E. Rains for drawing to our attention possible connections with [11]. This work was partially supported by EPSRC, The Leverhulme Trust, DFG, the European Union EQUIP project and the ESF QIT program, and Grants No. MEC (AP99), No. IUAP-P4-02, and No. GOA-Mefisto-666.

[1] M. B. Plenio and V. Vedral, *Contemp. Phys.* **39**, 431 (1998).

[2] C. H. Bennett *et al.*, *Phys. Rev. A* **53**, 2046 (1996); M. A. Nielsen, *Phys. Rev. Lett.* **83**, 436 (1999); G. Vidal, *ibid.* **83**, 1046 (1999); D. Jonathan and M. B. Plenio, *ibid.* **83**, 1455

(1999); L. Hardy, *Phys. Rev. A* **60**, 1912 (1999); H.-K. Lo and S. Popescu, e-print quant-ph/9707038.

[3] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, *Phys. Rev. A* **54**, 3824 (1996).

[4] E. M. Rains, *Phys. Rev. A* **60**, 173 (1999); **60**, 179 (1999).

[5] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, *Phys. Rev. Lett.* **78**, 2275 (1997); V. Vedral and M. B. Plenio, *Phys. Rev. A* **57**, 1619 (1998); V. Vedral, M. B. Plenio, K. A. Jacobs, and P. L. Knight, *Phys. Rev. A* **56**, 4452 (1997); M. B. Plenio, S. Virmani, and P. Papadopoulos, *J. Phys. A* **33**, L193 (2000); M. B. Plenio and V. Vedral, *J. Phys. A* **34**, 6997 (2001).

[6] For all subadditive measures of entanglement E one can show that $\limsup_{n \rightarrow \infty} E(\sigma^{\otimes n})/n = \lim_{n \rightarrow \infty} E(\sigma^{\otimes n})/n$; see H. Barnum, M. A. Nielsen, and B. Schumacher, *Phys. Rev. A* **57**, 4153 (1998).

[7] P. M. Hayden *et al.*, e-print quant-ph/0008134.

[8] If $E(\sigma^{\otimes n}) = nE(\sigma)$ for all $n = 1, 2, \dots$ for a given state σ , then clearly $E^\infty(\sigma) = E(\sigma)$ holds.

[9] R. F. Werner, *Phys. Rev. A* **40**, 4277 (1989).

[10] M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **84**, 4260 (2000). Despite many results of this paper being withdrawn, the formula for the Werner states' logarithm of the negativity is correct.

[11] E. Rains, e-print quant-ph/0008047.

[12] K. G. H. Vollbrecht and R. F. Werner, e-print quant-ph/0010095.

[13] A rigorous argument is as follows. Denote the maximum absolute value of $\partial \xi(k)/\partial k$ over $(0, n)$ by D . Then the error


$$\epsilon = \left| \sum_{k=0}^n C_n^k p^{n-k} (1-p)^k \xi(n-k) - \xi(pn) \right| \leq D \sum_{k=0}^n C_n^k p^{n-k} (1-p)^k |n-k-pn|,$$

because $\xi(n-k)$ lies between the bounds $\xi(pn) \pm D|n-k-pn|$. The last term is proportional to the mean absolute deviation from the mean, which is always less than the standard deviation (this follows from the concavity of $f(b) = |\sqrt{b}|$). Hence $\epsilon \leq D\sqrt{np(1-p)}$. In the case at hand, D is of order $1/n$: from the definition of $\xi(k)$, from Eq. (9) the maximal absolute value of $\partial \xi(k)/\partial k$ equals $\max_i |\lg(u_i/v_i)|/n$, and, as can be seen from Eq. (12), the values of u_i and v_i are independent of n . Hence, in the limit $n \rightarrow \infty$, the error ϵ goes to zero as $1/\sqrt{n}$.

[14] S. Boyd and L. Vandenberghe, *Introduction to Convex Optimization*, Lecture Notes (Stanford University, Stanford, CA, 2000/2001).

[15] The twirling operation Π in $d \times d$ dimensions, $d > 2$, is defined as $\Pi(\rho) = \int d\mu_U (U \otimes U) \rho (U \otimes U)^\dagger$ for states ρ , where the integral is performed with respect to the Haar measure, $U \in U(d)$. Under twirling, Werner states on $\mathbb{C}^2 \otimes \mathbb{C}^2$ with weight p of the antisymmetric state—regarded as states on $\mathbb{C}^d \otimes \mathbb{C}^d$ —will be mapped on Werner states on $\mathbb{C}^d \otimes \mathbb{C}^d$ with the same weight p .

[16] B. M. Terhal and K. G. H. Vollbrecht, *Phys. Rev. Lett.* **85**, 2625 (2000).



Simulated free energy of a 200-monomer chain as a function of crystal thickness (L) and orientational order parameter (S), dictating spontaneous selection of polymer crystal thickness. Ordered (valley) and disordered (saddle) configurations are shown. See article 218302.

PHYSICAL REVIEW LETTERS

Contents

VOLUME 87, NUMBER 21

19 November 2001

General Physics

Bose-Fermi Variational Theory of the Bose-Einstein Condensate Crossover to the Tonks Gas 210401
 M. D. Girardeau and E. M. Wright

Vortex Nucleation in a Stirred Bose-Einstein Condensate 210402
 C. Raman, J. R. Abo-Shaer, J. M. Vogels, K. Xu, and W. Ketterle

Driving Bose-Einstein-Condensate Vorticity with a Rotating Normal Cloud 210403
 P. C. Haljan, I. Coddington, P. Engels, and E. A. Cornell

Classical-Field Method for Time Dependent Bose-Einstein Condensed Gases 210404
 Alice Sinatra, Carlos Lobo, and Yvan Castin

Solution of the Relativistic Dirac-Morse Problem 210405
 A. D. Alhaidari

Collapse in Systems with Attractive Nonintegrable Potentials 210601
 I. Ispolatov and E. G. D. Cohen

Elementary Particles and Fields

Geometric CP Violation with Extra Dimensions 211601
 Darwin Chang and R. N. Mohapatra

Nonlinear Micromechanical Casimir Oscillator 211801
 H. B. Chan, V. A. Aksyuk, R. N. Kleiman, D. J. Bishop, and Federico Capasso

Bloch-Nordsieck Violation in Spontaneously Broken Abelian Theories 211802
 Marcello Ciafaloni, Paolo Ciafaloni, and Denis Comelli

Supersymmetric $SO(10)$ Grand Unified Models with Yukawa Unification and a Positive μ Term 211803
 Howard Baer and Javier Ferrandis

Charged-Particle Multiplicity $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV 211804
 T. Affolder *et al.* (CDF Collaboration)

Nuclear Physics

Energy Dependence of Nuclear Transparency in $C(p, 2p)$ Scattering 212301
 A. Leksanov, J. Alster, G. Asryan, Y. Averichev, D. Barton, V. Baturin, N. Bukhtoyarova, A. Carroll, S. Heppelmann, T. Kawabata, Y. Makdissi, A. Malki, E. Minina, I. Navon, H. Nicholson, A. Ogawa, Yu. Panebratsev, E. Piassetzky, A. Schetkovsky, S. Shimanskiy, A. Tang, J. W. Watson, H. Yoshida, and D. Zhalov

Effective Charge of the $\pi h_{11/2}$ Orbital and the Electric Field Gradient of Hg from the Yrast Structure of ^{206}Hg 212501
 B. Fornal, R. Broda, K. H. Maier, J. Wrzesiński, G. J. Lane, M. Cromaz, A. O. Macchiavelli, R. M. Clark, K. Vetter, A. P. Byrne, G. D. Dracoulis, M. P. Carpenter, R. V. F. Janssens, I. Wiedenhoever, M. Rejmund, and J. Blomqvist

Observation of a ${}^6_{\Lambda\Lambda}\text{He}$ Double Hypernucleus 212502
 H. Takahashi *et al.*

(Continued on Third Cover)

PRL 87 (21), 210401–219802, 19 November 2001 (256 total pages)

Contents (Continued)

Atomic, Molecular, and Optical Physics

- Generation of Electromagnetic Pulses from Plasma Channels Induced by Femtosecond Light Strings 213001
Chung-Chieh Cheng, E.M. Wright, and J.V. Moloney
- Benchmark Nonperturbative Calculations for the Electron-Impact Ionization of Li(2s) and Li(2p) 213201
J. Colgan, M.S. Pindzola, D.M. Mitnik, D.C. Griffin, and I. Bray

Nonlinear Dynamics, Fluid Dynamics, Classical Optics, etc.

- Stochastic Resonances in an Optical Two-Order Parameter Vectorial System 213901
Kamal P. Singh, Guy Ropars, Marc Brunel, Fabien Bretenaker, and Albert Le Floch
- Self-Guided Propagation of Ultrashort IR Laser Pulses in Fused Silica 213902
S. Tzortzakis, L. Sudrie, M. Franco, B. Prade, A. Mysyrowicz, A. Couairon, and L. Bergé
- Measurement of Lagrangian Velocity in Fully Developed Turbulence 214501
N. Mordant, P. Metz, O. Michel, and J.-F. Pinton
- Exact Calculation of the Linear Term in the Density Expansion of the Dynamic Structure Factor of a Dilute Gas ... 214502
Kunimasa Miyazaki and I.M. de Schepper

Plasma and Beam Physics

- Fast Proton Generation from Ultrashort Laser Pulse Interaction with Double-Layer Foil Targets 215001
J. Badziak, E. Woryna, P. Parys, K. Yu. Platonov, S. Jabłoński, L. Ryć, A.B. Vankov, and J. Wołowski
- Explosive Vaporization of Metallic Sodium Microparticles by CW Resonant Laser Radiation 215002
S.N. Atutov, W. Baldini, V. Biancalana, R. Calabrese, V. Guidi, B. Mai, E. Mariotti, G. Mazzocca,
L. Moi, S.P. Pod'yachev, and L. Tomassetti
- Finite Larmor-Radius Theory of Magnetic Island Evolution 215003
F.L. Waelbroeck, J.W. Connor, and H.R. Wilson
- Measurements of the Inverse Faraday Effect from Relativistic Laser Interactions with an Underdense Plasma 215004
Z. Najmudin, M. Tatarakis, A. Pukhov, E.L. Clark, R.J. Clarke, A.E. Dangor, J. Faure, V. Malka,
D. Neely, M.I.K. Santala, and K. Krushelnick

Condensed Matter: Structure, etc.

- Transition from Cage Clathrate to Filled Ice: The Structure of Methane Hydrate III 215501
J.S. Loveday, R.J. Nelmes, M. Guthrie, D.D. Klug, and J.S. Tse
- Thermal Transport Measurements of Individual Multiwalled Nanotubes 215502
P. Kim, L. Shi, A. Majumdar, and P.L. McEuen
- Model for a Random Laser 215503
A.L. Burin, M.A. Ratner, H. Cao, and R.P.H. Chang
- Neutron Investigation of the Ion Dynamics in Liquid Mercury: Evidence for Collective Excitations 215504
L.E. Bove, F. Sacchetti, C. Petrillo, B. Dörner, F. Formisano, and F. Barocchi
- Correlation of Positron Annihilation and Other Dynamic Properties in Small Molecule Glass-Forming Substances ... 215901
Kia L. Ngai, Li-Rong Bao, Albert F. Yee, and Christopher L. Soles
- Casimir Effect in Fluids above the Isotropic-Lamellar Transition 216101
Nariya Uchida
- Light-Atom Location in Adsorbed Benzene by Experiment and Theory 216102
G. Held, W. Braun, H.-P. Steinrück, S. Yamagishi, S.J. Jenkins, and D.A. King
- Kinetics of Faceting Driven by Attractive Step-Step Interactions on Vicinal Si(113) 216103
K. Sudoh and H. Iwasaki

Condensed Matter: Electronic Properties, etc.

- Fermi Surface Nesting in Disordered Cu_{1-x}Pd_x Alloys 216401
I. Wilkinson, R.J. Hughes, Zs. Major, S.B. Dugdale, M.A. Alam, E. Bruno, B. Ginatempo,
and E.S. Giuliano
- Charged Excitons in the Fractional Quantum Hall Regime 216402
G. Yusa, H. Shtrikman, and I. Bar-Joseph
- Muonium Analog of Hydrogen Passivation: Observation of the Mu⁺-Zn⁻ Reaction in GaAs 216403
K.H. Chow, B. Hitti, R.F. Kiefl, R.L. Lichti, and T.L. Estle

(Continued on Preceding Page)

Contents (Continued)

Spectroscopic Indications of Polaronic Carriers in the Quasi-One-Dimensional Conductor $(\text{TaSe}_4)_2\text{I}$	216404
L. Perfetti, H. Berger, A. Reglinelli, L. Degiorgi, H. Höchst, J. Voit, G. Margaritondo, and M. Gioni	
Importance of Correlation Effects on Magnetic Anisotropy in Fe and Ni	216405
Imseok Yang, Sergej Y. Savrasov, and Gabriel Kotliar	
Kondo Effect in Real Quantum Dots	216601
M. Pustilnik and L.I. Glazman	
Resistance Spikes and Domain Wall Loops in Ising Quantum Hall Ferromagnets	216801
T. Jungwirth and A.H. MacDonald	
Metallic Phase in Quantum Hall Systems due to Inter-Landau-Band Mixing	216802
Gang Xiong, Shi-Dong Wang, Qian Niu, De-Cheng Tian, and X.R. Wang	
Tunneling through a Quantum Dot in Local Spin Singlet-Triplet Crossover Region with Kondo Effect	216803
Wataru Izumida, Osamu Sakai, and Seigo Tarucha	
Observation of Magnetically Induced Effective-Mass Enhancement of Quasi-2D Excitons	216804
L.V. Butov, C.W. Lai, D.S. Chemla, Yu.E. Lozovik, K.L. Campman, and A.C. Gossard	
Crossover from Electronic to Atomic Shell Structure in Alkali Metal Nanowires	216805
A.I. Yanson, I.K. Yanson, and J.M. van Ruitenbeek	
Photoinduced Anisotropy of Photoconductivity in Amorphous $\text{As}_{50}\text{Se}_{50}$ Films	216806
V. Lyubin, M. Klebanov, and V.K. Tikhomirov	
Shot-Noise in Transport and Beam Experiments	216807
U. Gavish, Y. Levinson, and Y. Imry	
Resonant Transport in Nb/GaAs/AlGaAs Heterostructures: Realization of the de Gennes–Saint-James Model	216808
Francesco Giazotto, Pasqualantonio Pingue, Fabio Beltram, Marco Lazzarino, Daniela Orani, Silvia Rubini, and Alfonso Franciosi	
Doping Dependence of Pseudogap and Related Charge Dynamics in $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$	217001
Y. Onose, Y. Taguchi, K. Ishizaka, and Y. Tokura	
Projected Wave Functions and High Temperature Superconductivity	217002
Arun Paramakanti, Mohit Randeria, and Nandini Trivedi	
Quantum Phase Slips in Superconducting Nanowires	217003
C.N. Lau, N. Markovic, M. Bockrath, A. Bezryadin, and M. Tinkham	
Quantized Thermal Hall Effect in the Mixed State of d -Wave Superconductors	217004
Ashvin Vishwanath	
Spectacular Doping Dependence of Interlayer Exchange and Other Results on Spin Waves in Bilayer Manganites ...	217201
T.G. Perring, D.T. Adroja, G. Chaboussant, G. Aeppli, T. Kimura, and Y. Tokura	
Orbital Excitations in LaMnO_3	217202
Jeroen van den Brink	
Spin-Wave Instabilities in Large-Scale Nonlinear Magnetization Dynamics	217203
Giorgio Bertotti, Isaak D. Mayergoyz, and Claudio Serpico	
Gilbert Damping in Single and Multilayer Ultrathin Films: Role of Interfaces in Nonlocal Spin Dynamics	217204
R. Urban, G. Woltersdorf, and B. Heinrich	
Hopping Transport of Positrons in Hydrogenated Amorphous Silicon	217401
D.T. Britton, A. Hempel, and W. Triftshäuser	
Evidence for Pb-O Covalency in Tetragonal PbTiO_3	217601
Yoshihiro Kuroiwa, Shinobu Aoyagi, Akikatsu Sawada, Jimpei Harada, Eiji Nishibori, Masaki Takata, and Makoto Sakata	
Interdisciplinary Physics: Biological Physics, Quantum Information, etc.	
Quantum Solution to the Byzantine Agreement Problem	217901
Matthias Fitzl, Nicolas Gisin, and Ueli Maurer	
Asymptotic Relative Entropy of Entanglement	217902
K. Audenaert, J. Eisert, E. Jané, M.B. Plenio, S. Virmani, and B. De Moor	
Positional, Reorientational, and Bond Orientational Order in DNA Mesophases	218101
V. Lorman, R. Podgornik, and B. Žekš	
Nonstationary Phenomena in Tapered Gyro-Backward-Wave Oscillators	218301
G.S. Nusinovich, A.N. Vlasov, and T.M. Antonsen, Jr.	

(Continued on Preceding Page)

Contents (Continued)

Molecular Mechanisms of Polymer Crystallization from Solution	218302
P. Welch and M. Muthukumar	
Comments	
Superfluidity at Supersonic Speed?	218901
Mourad Haddad and Vincent Hakim	
Law <i>et al.</i> Reply	218902
C. K. Law, C. M. Chan, P. T. Leung, and M. C. Chu	
Comment on "Breakdown of the Internet under Intentional Attack"	219801
S. N. Dorogovtsev and J. F. F. Mendes	
Cohen, Erez, ben-Avraham, and Havlin Reply	219802
Reuven Cohen, Keren Erez, Daniel ben-Avraham, and Shlomo Havlin	