Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

http://www.elsevier.com/copyright

Author's personal copy

Biochimica et Biophysica Acta 1798 (2010) 1908–1912

Contents lists available at ScienceDirect

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbamem

YTPdb: A wiki database of yeast membrane transporters

Sylvain Brohée [a,1], Roland Barriot [a,b,c,2], Yves Moreau [a,3], Bruno André [d,*]

[a] ESAT-SCD, Katholieke Universiteit Leuven, Kasteelpark Arenberg 10, B-3001 Leuven, Belgium
[b] Université de Toulouse, UPS, Laboratoire de Microbiologie et Génétique Moléculaires, F-31000 Toulouse, France
[c] Centre National de la Recherche Scientifique, LMGM, F-31000 Toulouse, France
[d] Physiologie Moléculaire de la Cellule, Institut de Biologie et de Médecine Moléculaires, Université Libre de Bruxelles, Rue des Pr, Jeener et Brachet, B-6041 Gosselies, Belgium

ARTICLE INFO

Article history:
Received 19 February 2010
Received in revised form 17 May 2010
Accepted 7 June 2010
Available online 19 June 2010

Keywords:
yeast
transporter
database

ABSTRACT

Membrane transporters constitute one of the largest functional categories of proteins in all organisms. In the yeast *Saccharomyces cerevisiae*, this represents about 300 proteins (~5% of the proteome). We here present the Yeast Transport Protein database (YTPdb), a user-friendly collaborative resource dedicated to the precise classification and annotation of yeast transporters. YTPdb exploits an evolution of the MediaWiki web engine used for popular collaborative databases like Wikipedia, allowing every registered user to edit the data in a user-friendly manner. Proteins in YTPdb are classified on the basis of functional criteria such as subcellular location or their substrate compounds. These classifications are hierarchical, allowing queries to be performed at various levels, from highly specific (e.g. ammonium as a substrate or the vacuole as a location) to broader (e.g. cation as a substrate or inner membranes as location). Other resources accessible for each transporter via YTPdb include post-translational modifications, K_m values, a permanently updated bibliography, and a hierarchical classification into families. The YTPdb concept can be extrapolated to other organisms and could even be applied for other functional categories of proteins. YTPdb is accessible at http://homes.esat.kuleuven.be/ytpdb/.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

The *membrane transporter* functional class of proteins is typically one of the most abundantly represented in cellular proteomes. The estimated percentage of predicted membrane transporters ranges from 2% to 10% in bacteria and 2% to 7% in eukaryotes. It is 2.7% in the human species [1]. These proteins facilitate the transport of a very wide variety of small compounds (small ions, organic compounds, short peptides, water, etc.) across the lipid bilayers of the plasma and internal membranes. Most inventoried transporters belong to families conserved throughout evolution [2].

Among the 5690 protein-encoding genes of the yeast *Saccharomyces cerevisiae* [3], almost 300 code for established or predicted transmembrane transporters [4]. *Established* transporters are those for which experimental evidence of a direct role in transmembrane transport of specific compounds is available. *Predicted* transporters share significant sequence similarity with at least one experimentally characterized transporter from any organism, but their functions remain unknown: although the subcellular locations of many of them have been determined by large-scale analysis [5], the compounds they recognize have not been identified and their roles in cell metabolism thus remain unknown.

Specialized databases of membrane transporters have previously been implemented (links accessible via YTPdb). For instance, the Transport Classification Database (TCDB) provides a classification system for membrane transporters of all organisms into no less than 550 families [2], and the related TransportDB lists all transporters predicted from completely sequenced genomes [1]. Other databases categorize transporters or membrane proteins of a single or a small number of organisms, e.g. the Yeast Transport Information (YETI) [6], the Yeast Membrane Protein Library (YMPL) [7], or the Plant Membrane protein Database (Aramemnon) [8]. There also exist databases supplying information about one type of transporters in different organisms (e.g. P-type ATPases [9]). Lastly, information about membrane transporters is also accessible through databases collecting data on the complete sets of genes and proteins of specific organisms, e.g. the Saccharomyces Genome Database (SGD) [10] or the Comprehensive Yeast Genome Database (CYGD) [11].

Yet molecular biologists working in the membrane transport field also need databases providing classification systems based on the main functional properties of listed proteins, e.g. their substrate compounds and subcellular locations. Furthermore, these databases should ideally include annotations about the affinity constants (K_m), membrane topology, post-translational modifications, as well as an

* Corresponding author. Tel.: +32 2 6509958.
E-mail addresses: Sylvain.Brohee@esat.kuleuven.be (S. Brohée), roland.barriot@biotoul.fr (R. Barriot), yves.moreau@esat.kuleuven.be (Y. Moreau), bran@ulb.ac.be (B. André).
[1] Tel.: +32 16 328643.
[2] Tel.: +33 5 61335821.
[3] Tel.: +32 16 328645.

0005-2736/$ – see front matter © 2010 Elsevier B.V. All rights reserved.
doi:10.1016/j.bbamem.2010.06.008

Author's personal copy

up-to-date bibliography. It seems obvious that such goal is only achievable if this database can be updated through a user-friendly interface by several users having expertise in the field. We have applied these criteria to implement the Yeast Transport Protein database (YTPdb: http://homes.esat.kuleuven.be/ytpdb/), a web-accessible database devoted to the precise classification and annotation of *S. cerevisiae* membrane transport proteins. Besides its main features described below, YTPdb is presented in the form of a wiki, i.e. on the model of the well-known web encyclopedia Wikipedia. Any registered user can thus freely edit data in the database and add or remove data. We think that the YTPdb concept is extensible to other organisms and even to other functional classes of proteins.

2. Results

2.1. YTPdb content

The Yeast Transporter Protein database provides manually annotated information on 299 yeast proteins classified as *established* (199) or *predicted* (100) membrane transporters (see Introduction). These 299 proteins correspond to the 337 proteins predicted to be transporters by the TransportTP server [12] from which 38 have not been considered because they correspond to subunits of protein translocation systems, of the mitochondrial ATP synthase, of the vacuolar H^+ pump, to non-membrane proteins containing the ABC (ATP-binding cassette) motif, and to several enzymes (e.g. HMG-CoA reductases, lipid modification enzymes, etc.) and membrane proteins which may not be classified as *established* or *predicted* transporters. For each protein, YTPdb provides a page including a short description, the list of substrates, and the subcellular location (Fig. 1). Clicking on a substrate gives the list of all proteins transporting this compound and clicking on a protein's location gives the list of proteins colocalized in this cell compartment. An originality of YTPdb is that functional classification criteria such as subcellular location and recognized substrates are organized in a tree-like fashion. For instance, a *glucose* transporter is also classified as a transporter of *hexose*, *monosaccharide*, and *carbohydrate* and a *cytosine* transporter as a protein transporting *pyrimidine* and *nucleobase*. Similarly, a protein localizing to the *Golgi compartment* is also classified as a protein present in the *Golgi and endosome membrane system*, in the *secretory pathway membrane system*, and in *internal membranes*. It is thus possible – using the "Browse by substrate", "Browse by substrate categories", and "Browse by subcellular location" query tools – to retrieve proteins on the basis of very specific or broader criteria. For instance, the user can list all calcium or leucine transporters but also those for cations, amino acids, etc. Similarly, the user can view all transporters present in the "late endosome" or in the less defined "Golgi and endosomal membrane" system. This mode of classification is convenient in the case of proteins whose properties have been characterized with only limited precision. For instance, the P-type ATPase Neo1 has been located in the Golgi and/or endosomal membrane [13,14]. YTPdb contains a total list of 315 chemical compounds (plus 65 synonyms) and 28 subcellular locations, corresponding to the branch extremities of the tree-like classification schemes. The affinity constants (K_m) for substrates (136 values in the May 2010 version of YTPdb) with the corresponding references are also accessible.

Transporters encoded in YTPdb are also classified according to the transport classification (TC) system [2] (Fig. 1). Using the "Browse by TC class" option, proteins at any level of the TC classification can be listed, e.g. all *channels/pores* (22 proteins) or the *Major Intrinsic Protein (MIP) Family* (4 proteins). We have developed another classification system (YTPdb classification) including more subdivisions and more suited to yeast transporters. For instance, P-type ATPases (16 proteins in *S. cerevisiae*) form a unique class of proteins in TC, whereas they are subdivided into six subfamilies in the YTP classification. The YTP classification also provides a multiple sequence alignment computed by the Muscle algorithm [15].

An important feature of transporters is the topology they adopt in the membrane. For each protein, YTPdb gives access to a web page recapitulating the data of topology prediction generated by seven distinct algorithms.

Many protein databases do not provide an updated bibliography for protein entries. In YTPdb, two systems have been implemented to associate each transporter with references. The "All references" option is based on an editable Boolean formula retrieving references from the PubMed.gov server, and the corresponding bibliography is thus permanently updated. Through the "Curated references" option, the user can generate a list of references classified into several categories. For instance, "Early studies" provides references about studies preceding the molecular characterization of the transporter gene, "Initial molecular characterization, general properties and function" include references about the functional characterisation of the transporter. Other categories (eight in all) list references about "Intracellular trafficking and its regulation", "Gene expression and its regulation", "Transport activity (including influence of mutations) and its regulation", etc. Curated lists of references are currently available for about 40 transporters. As mentioned below, registered curators are invited to associate references with their favorite transporters.

Post-translational modifications (PTMs) play a central role in regulation of protein function. This is well illustrated for yeast transporters, e.g., ubiquitylation typically targets these proteins for degradation in the vacuole [16] and many yeast transporters have been shown to be phosphorylated. A special effort has been devoted to including in YTPdb all known PTMs of yeast transporters deduced from studies centered on specific proteins and from large-scale mass spectrometry analyses [17–25]. In the current version of YTPdb (May 2010), 764 unique PTMs have been encoded for 123 transporters. For each protein, these PTMs are presented in a recapitulative table with references, and their relative positions are indicated in a figure showing the predicted transmembrane regions [26], global topology, and positions of conserved PFAM domains [27] of the protein (Fig. 2). This convenient presentation facilitates comparisons between proteins of the same family and detection of possible recurrent PTM profiles. For instance, many amino acid permeases are palmitoylated at their C-termini [28], and their cytosolic N-terminal tails often contain several phosphorylated serines and threonines as well as ubiquitylated lysines.

YTPdb also associates each transporter with direct links to well-known databases (MIPS [29], Ensembl [30], SGD [10]) to a repository of gene expression data (ArrayExpress) [31] and to the Yeast Gene Order Browser [32]. The latter is an online tool for visualizing the syntenic context of any gene from twelve hemiascomycete species; this is particularly useful for detecting the many pairs of transporter genes deriving from the whole genome duplication event that occurred in an ancestor of the *S. cerevisiae* lineage [33,34].

2.2. YTPdb is a collaborative database

There is no longer any need to introduce Wiki technology. The past few years have seen an explosion in the number of wiki-based web sites, ranging from very general (e.g. the well-known Wikipedia, wiktionnary) to more specialized or biological (wikiproteins [35], wikipathways [36], wikipedia for genes [37], etc). YTPdb is a wiki based on an extension of MediaWiki technology (running Wikipedia). This implies that almost all data in the database can be freely edited and extended by any registered user. For instance, specific and readily editable forms are available for adding K_m values, post-translational modifications, and references and for editing the basic information (substrate compounds, subcellular location, gene synonyms, Boolean formula to retrieve references, etc.) on each transporter. A similar form is available for adding or editing substrate compounds

Author's personal copy



Fig. 1. A YTPdb protein report. The "Family" section gives access to the protein family and upper-level families to which the protein belongs; the "Substrates" section presents the list of compounds recognized by the transporter; the "Location" section gives access to its subcellular location. By clicking on a given substrate or on the subcellular location, all transporters recognizing this compound or colocalized with the protein are shown. The other sections provide access to a list of K_m values for different substrates, post-translational modifications, external links, sequences, and references (see text). Numbers between brackets specify the quantity of encoded data. By clicking on "edit" links, the user can introduce free text (e.g. Remarks).

(synonyms, classification). Furthermore, by clicking on the "Edit" link at the top of a page (or on the "Create" link, when free text has not yet been included in the page), it is possible to include additional comments and figures about, e.g., a specific transporter (Fig. 1), transporter family, substrate compound, cell compartment, etc. For this, the user is also redirected to a form where free text can be entered as in standard word processors. Finally, as in Wikipedia, it is possible to create from scratch new pages editable by users. Extensive information about the editing process of YTP is available in the Tutorial section of the database.

One drawback of classical wikis is that the data are stored in free text format in a main unique underlying database. This differs from

Author's personal copy



Post translational modifications of protein Gap1

Add a new post-tranlational modification

Position	Modification	Comment	References
K 9	ubiquitylation	K63-linked ubiquitylation	▪ Soetens O *et al* (2001) ▪ Lauwers E *et al* (2009)
K 16	ubiquitylation	K63-linked ubiquitylation	▪ Soetens O *et al* (2001) ▪ Lauwers E *et al* (2009)
K 76	ubiquitylation		▪ Hitchcock AL *et al* (2003) ▪ Peng J *et al* (2003)
C 602	palmitoylation		▪ Roth AF *et al* (2006)

Relative position of modified residues in Gap1

Transmembrane segments (upper figure) were predicted using TMHMM.

Conserved functional domains (lower figure) were predicted using Interpro - PFAM.

Fig. 2. Example of a report presenting the post-translational modifications associated with a specific yeast transporter. These modifications are reported on a protein map, together with the PFAM functional domains and the predicted transmembrane segments.

most biological databases, where data are stored in well-structured repositories, and web pages are generated by extracting a set of entries from different tables. To circumvent this problem, YTPdb makes use of *WikiOpener*, an extension for the classical MediaWiki engine that augments Wiki pages by allowing on-the-fly querying, editing, and formatting resources external to the Wiki (MediaWiki web site: http://www.mediawiki.org/wiki/Extension:WikiOpener). Those resources may correspond to data extracted from databases, flat files, DAS tracks, or even results returned by local or remote bioinformatics analysis tools. WikiOpener has already been successfully used in the frame of another biological database that aims at depicting the relationships between human heart defects and gene alterations [38]. In the case of YTPdb, *WikiOpener* mainly queries a relational database, distinct from the main database used to populate the pages of the wiki. Moreover, WikiOpener uses the PubMed identifier of references to connect to the NCBI web services and gather the references details which are stored in the YTPdb relational database. WikiOpener also determines the presentation format of these queries within the wiki. The PHP code of the YTPdb components as well as the scheme of the relational database can be obtained by simple request to the authors.

3. Discussion

YTPdb displays several features distinguishing it from other protein databases and useful to experts in the transport field as well as to occasional users. Transporters can be retrieved from hierarchical classification systems based on substrate specificity, subcellular location, and similarities of primary structure. YTPdb also includes K_m values and all reported post-translational modifications. Importantly, these annotations are based solely on experimental data available in the literature and source references are indicated. Furthermore, YTPdb is a collaborative database, meaning that any registered user is invited to modify and update entries related to his/her expertise. The web engine running YTPdb is the same as the one running the Wikipedia web encyclopedia, making editing and adding information easy and straightforward for people having less computer skills. The advantage of a collaborative database is that it can be constantly updated by multiple curators. This reduces the risk of the database progressively becoming obsolete, as often occurs with small specialized biological databases [39–41]. The disadvantage of a collaborative database is that users can potentially add wrong information. In the case of YTPdb, however, curators have to register (a difference with Wikipedia) allowing the administrator to limit access of YTPdb to scientists active in the field. We propose that the concept and structure of YTPdb is transposable to other species for which availability of a transporter database would be beneficial to the scientific community. We also think that YTPdb can serve as a model for databases of other functional categories of proteins.

Acknowledgements

Many thanks to Daniel Van Belle for creating with BA the initial version of YTPdb. We thank Jacques van Helden, Robert Herzog, Marc Colet, and Antonio Urrestarazu for constant support and encouragement. We also thank Morgane Thomas-Chollier, Elsa Lauwers, Eckhard Boles, and Valérie Ledent for many helpful discussions. Finally, we thank Olivier Cagnac, Hana Sychrova, Emmanuel Lesuisse, and Juergen Stolz for adding the first external contributions to YTPdb. SB was the recipient of a PhD grant from the Belgian Fonds pour la

Author's personal copy

Formation à la Recherche dans l'Industrie et l'Agriculture (FRIA) and is now supported by the E.U. FP7 CHeartED project. KULeuven SCD-SISTA lab is funded by grants provided by the (1) Research Council KUL: GOA MaNet, CoE EF/05/007 SymBioSys; (2) IWT: SBO-MoKa; (3) the Belgian Federal Science Policy Office: IUAP P6/25 (BioMaGNet); and (4) the European Union: FP7-HEALTH CHeartED.

References

[1] Q. Ren, K. Chen, I.T. Paulsen, TransportDB: a comprehensive database resource for cytoplasmic membrane transport systems and outer membrane channels, Nucleic Acids Res. 35 (2007) D274–D279.
[2] M.H. Saier Jr., M.R. Yen, K. Noto, D.G. Tamang, C. Elkan, The Transporter Classification Database: recent advances, Nucleic Acids Res. 37 (2009) D274–D278.
[3] A. Goffeau, B.G. Barrell, H. Bussey, R.W. Davis, B. Dujon, H. Feldmann, F. Galibert, J.D. Hoheisel, C. Jacq, M. Johnston, E.J. Louis, H.W. Mewes, Y. Murakami, P. Philippsen, H. Tettelin, S.G. Oliver, Life with 6000 genes, Science 274 (1996) 546, 563–546, 567.
[4] D. Van Belle, B. André, A genomic view of yeast membrane transporters, Curr. Opin. Cell Biol. 13 (2001) 389–398.
[5] W.K. Huh, J.V. Falvo, L.C. Gerke, A.S. Carroll, R.W. Howson, J.S. Weissman, E.K. O'Shea, Global analysis of protein localization in budding yeast, Nature 425 (2003) 686–691.
[6] B. De Hertogh, E. Carvajal, E. Talla, B. Dujon, P. Baret, A. Goffeau, Phylogenetic classification of transporters and other membrane proteins from *Saccharomyces cerevisiae*, Funct. Integr Genomics 2 (2002) 154–170.
[7] J.M. Ward, Identification of novel families of membrane proteins from the model plant *Arabidopsis thaliana*, Bioinformatics. 17 (2001) 560–563.
[8] R. Schwacke, A. Schneider, E. van der Graaff, K. Fischer, E. Catoni, M. Desimone, W.B. Frommer, U.I. Flugge, R. Kunze, ARAMEMNON, a novel database for Arabidopsis integral membrane proteins, Plant Physiol 131 (2003) 16–26.
[9] K.B. Axelsen, M.G. Palmgren, Evolution of substrate specificities in the P-type ATPase superfamily, J. Mol. Evol. 46 (1998) 84–101.
[10] S.R. Engel, R. Balakrishnan, G. Binkley, K.R. Christie, M.C. Costanzo, S.S. Dwight, D.G. Fisk, J.E. Hirschman, B.C. Hitz, E.L. Hong, C.J. Krieger, M.S. Livstone, S.R. Miyasato, R. Nash, R. Oughtred, J. Park, M.S. Skrzypek, S. Weng, E.D. Wong, K. Dolinski, D. Botstein, J.M. Cherry, Saccharomyces Genome Database provides mutant phenotype data, Nucleic Acids Res. 38 (2010) D433–D436.
[11] U. Guldener, M. Munsterkotter, G. Kastenmuller, N. Strack, J. van Helden, C. Lemer, J. Richelle, S.J. Wodak, J. Garcia-Martinez, J.E. Perez-Ortin, H. Michael, A. Kaps, E. Talla, B. Dujon, B. Andre, J.L. Souciet, J. De Montigny, E. Bon, C. Gaillardin, H.W. Mewes, CYGD: the Comprehensive Yeast Genome Database, Nucleic Acids Res. 33 (2005) D364–D368.
[12] H. Li, X. Dai, X. Zhao, A nearest neighbor approach for automated transporter prediction and categorization from protein sequences, Bioinformatics. 24 (2008) 1129–1136.
[13] S. Wicky, H. Schwarz, B. Singer-Kruger, Molecular interactions of yeast Neo1p, an essential member of the Drs2 family of aminophospholipid translocases, and its role in membrane trafficking within the endomembrane system, Mol. Cell Biol. 24 (2004) 7402–7418.
[14] Z. Hua, T.R. Graham, Requirement for neo1p in retrograde transport from the Golgi complex to the endoplasmic reticulum, Mol. Biol. Cell 14 (2003) 4971–4983.
[15] R.C. Edgar, MUSCLE: a multiple sequence alignment method with reduced time and space complexity, BMC, Bioinformatics 5 (2004) 113.
[16] E. Lauwers, Z. Erpapazoglou, R. Haguenauer-Tsapis, B. Andre, The ubiquitin code of yeast permease trafficking, Trends Cell Biol. 20 (2010) 196–204.
[17] A. Gruhler, J.V. Olsen, S. Mohammed, P. Mortensen, N.J. Faergeman, M. Mann, O.N. Jensen, Quantitative phosphoproteomics applied to the yeast pheromone signaling pathway, Mol. Cell Proteomics. 4 (2005) 310–327.
[18] X. Li, S.A. Gerber, A.D. Rudner, S.A. Beausoleil, W. Haas, J. Villen, J.E. Elias, S.P. Gygi, Large-scale phosphorylation analysis of alpha-factor-arrested *Saccharomyces cerevisiae*, J. Proteome. Res. 6 (2007) 1190–1197.
[19] A. Chi, C. Huttenhower, L.Y. Geer, J.J. Coon, J.E. Syka, D.L. Bai, J. Shabanowitz, D.J. Burke, O.G. Troyanskaya, D.F. Hunt, Analysis of phosphorylation sites on proteins from *Saccharomyces cerevisiae* by electron transfer dissociation (ETD) mass spectrometry, Proc. Natl. Acad. Sci. U.S.A. 104 (2007) 2193–2198.
[20] M.B. Smolka, C.P. Albuquerque, S.H. Chen, H. Zhou, Proteome-wide identification of in vivo targets of DNA damage checkpoint kinases, Proc. Natl. Acad. Sci. U.S.A. 104 (2007) 10364–10369.
[21] C.P. Albuquerque, M.B. Smolka, S.H. Payne, V. Bafna, J. Eng, H. Zhou, A multidimensional chromatography technology for in-depth phosphoproteome analysis, Mol. Cell Proteomics. 7 (2008) 1389–1396.
[22] A. Huber, B. Bodenmiller, A. Uotila, M. Stahl, S. Wanka, B. Gerrits, R. Aebersold, R. Loewith, Characterization of the rapamycin-sensitive phosphoproteome reveals that Sch9 is a central coordinator of protein synthesis, Genes Dev. 23 (2009) 1929–1943.
[23] A.L. Hitchcock, K. Auld, S.P. Gygi, P.A. Silver, A subset of membrane-associated proteins is ubiquitinated in response to mutations in the endoplasmic reticulum degradation machinery, Proc. Natl. Acad. Sci. U.S.A. 100 (2003) 12735–12740.
[24] J. Peng, D. Schwartz, J.E. Elias, C.C. Thoreen, D. Cheng, G. Marsischky, J. Roelofs, D. Finley, S.P. Gygi, A proteomics approach to understanding protein ubiquitination, Nat. Biotechnol. 21 (2003) 921–926.
[25] F. Gnad, L.M. de Godoy, J. Cox, N. Neuhauser, S. Ren, J.V. Olsen, M. Mann, High-accuracy identification and bioinformatic analysis of in vivo protein phosphorylation sites in yeast, Proteomics. 9 (2009) 4642–4652.
[26] A. Krogh, B. Larsson, Heijne G. von, E.L. Sonnhammer, Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes, J.Mol.Biol. 305 (2001) 567–580.
[27] S.J. Sammut, R.D. Finn, A. Bateman, Pfam 10 years on: 10, 000 families and still growing, Brief. Bioinform. 9 (2008) 210–219.
[28] A.F. Roth, J. Wan, A.O. Bailey, B. Sun, J.A. Kuchar, W.N. Green, B.S. Phinney, J.R. Yates III, N.G. Davis, Global analysis of protein palmitoylation in yeast, Cell 125 (2006) 1003–1013.
[29] H.W. Mewes, S. Dietmann, D. Frishman, R. Gregory, G. Mannhaupt, K.F. Mayer, M. Munsterkotter, A. Ruepp, M. Spannagl, V. Stumpflen, T. Rattei, MIPS: analysis and annotation of genome information in 2007, Nucleic Acids Res. 36 (2008) D196–D201.
[30] P. Flicek, B.L. Aken, B. Ballester, K. Beal, E. Bragin, S. Brent, Y. Chen, P. Clapham, G. Coates, S. Fairley, S. Fitzgerald, J. Fernandez-Banet, L. Gordon, S. Graf, S. Haider, M. Hammond, K. Howe, A. Jenkinson, N. Johnson, A. Kahari, D. Keefe, S. Keenan, R. Kinsella, F. Kokocinski, G. Koscielny, E. Kulesha, D. Lawson, I. Longden, T. Massingham, W. McLaren, K. Megy, B. Overduin, B. Pritchard, D. Rios, M. Ruffier, M. Schuster, G. Slater, D. Smedley, G. Spudich, Y.A. Tang, S. Trevanion, A. Vilella, J. Vogel, S. White, S.P. Wilder, A. Zadissa, E. Birney, F. Cunningham, I. Dunham, R. Durbin, X.M. Fernandez-Suarez, J. Herrero, T.J. Hubbard, A. Parker, G. Proctor, J. Smith, S.M. Searle, Ensembl's 10th year, Nucleic Acids Res. 38 (2010) D557–D562.
[31] M. Kapushesky, I. Emam, E. Holloway, P. Kurnosov, A. Zorin, J. Malone, G. Rustici, E. Williams, H. Parkinson, A. Brazma, Gene expression atlas at the European bioinformatics institute, Nucleic Acids Res. 38 (2010) D690–D698.
[32] K.P. Byrne, K.H. Wolfe, Visualizing syntenic relationships among the hemiascomycetes with the Yeast Gene Order Browser, Nucleic Acids Res. 34 (2006) D452–D455.
[33] K.H. Wolfe, D.C. Shields, Molecular evidence for an ancient duplication of the entire yeast genome, Nature 387 (1997) 708–713.
[34] M. Kellis, B.W. Birren, E.S. Lander, Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*, Nature 428 (2004) 617–624.
[35] B. Mons, M. Ashburner, C. Chichester, E. van Mulligen, M. Weeber, J. den Dunnen, G.J. van Ommen, M. Musen, M. Cockerill, H. Hermjakob, A. Mons, A. Packer, R. Pacheco, S. Lewis, A. Berkeley, W. Melton, N. Barris, J. Wales, G. Meijssen, E. Moeller, P.J. Roes, K. Borner, A. Bairoch, Calling on a million minds for community annotation in WikiProteins, Genome Biol 9 (2008) R89-.
[36] A.R. Pico, T. Kelder, M.P. van Iersel, K. Hanspers, B.R. Conklin, C. Evelo, WikiPathways: pathway editing for the people, PLoS.Biol. 6 (2008) e184-.
[37] J.W. Huss III, P. Lindenbaum, M. Martone, D. Roberts, A. Pizarro, F. Valafar, J.B. Hogenesch, A.I. Su, The Gene Wiki: community intelligence applied to human gene annotation, Nucleic Acids Res. 38 (2010) D633–D639.
[38] R. Barriot, J. Breckpot, B. Thienpont, S. Brohée, S. Van Vooren, B. Coessens, L.C. Tranchevent, P. Van Loo, M. Gewillig, K. Devriendt, Y. Moreau, Collaboratively charting the gene-to-phenotype network of human congenital heart defects, Genome Med. 2 (2010) 16.
[39] L.B. Ellis, D. Kalumbi, The demise of public data on the web? Nat. Biotechnol. 16 (1998) 1323–1324.
[40] L.B. Ellis, D. Kalumbi, Financing a future for public biological data, Bioinformatics. 15 (1999) 717–722.
[41] Chandras Christina, Weaver Thomas, Zouberakis Michae, Smedley Damian, Schughart Klaus, Rosenthal Nadia, Hancock John M, Kollias George, Schofield Paul N, Aidinis Vasilis, Models for financial sustainability of biological databases and resources, Databases 2009:bap017 (2009).