10 - 08

Elsevier Editorial System(tm) for Early Human Development
Manuscript Draft

Manuscript Number: EHD-D-09-00126R1

Title: Decoupling between fundamental frequency and energy envelope of neonate cries

Article Type: Research Paper

Keywords: cry analysis; decoupling; infant; maturation; Modified Behavioural Pain Scale

Corresponding Author: Prof. Daniel Berckmans, Ph.D., Ir.

Corresponding Author's Institution:

First Author: Mitchell Silva

Order of Authors: Mitchell Silva; Bogdan Mijovic; Bea R.H. Van den Bergh; Karel Allegaert; Jean-Marie Aerts; Sabine Van Huffel; Daniel Berckmans

Abstract: Background: The presence of decoupling, i.e. the absence of coupling between fundamental frequency variation and energy envelope during phonetic crying, and its extent, reflects the degree of maturation of the central nervous system.
Aim: We hereby wanted to assess the existence and extent of decoupling in term neonates (neurodevelopmental relevance) and whether an association between decoupling and clinical pain expression could be unveiled (clinical relevance).
Study design: To assess decoupling in healthy term neonates during procedural pain, newborns were videotaped and crying was recorded during venous blood sampling. Besides acoustic analysis, pain expression was quantified based on the Modified Behavioral Pain Scale (MBPS).
Subjects: 47 healthy term neonates underwent venous blood puncture at the 3rd day of life.
Outcome measures: Beside the MBPS-score, the correlation coefficients were calculated between the fundamental frequency variation and energy envelope of the cries.
Results: Based on data collected in 47 healthy term neonates, correlation coefficients varied between 0.20 and 0.68. The degree of decoupling displayed extensive variability between the neonates and also in different cry bouts in a crying sequence within an individual neonate. A negative association was found between MBPS-value and decoupling (r^2=-0.12), the same as for the intra-subject variability although less extensive (r^2=-0.02).
Conclusion: Decoupling only relates weakly with the amount of distress in 3 day old newborns, even though a great intra-subject variability is present. This study suggests that there is no evidence of extensive decoupling as the newborn still has to fully develop the control of larynx and abdominal muscles.

Cover Letter

FACULTY OF BIOSCIENCE ENGINEERING
DEPARTMENT OF BIOSYSTEMS
MEASURE, MODEL & MANAGE BIORESPONSES (M3-BIORES)
KASTEELPARK ARENBERG 30
B-3001 HEVERLEE (LEUVEN)
BELGIUM

KATHOLIEKE
UNIVERSITEIT
LEUVEN

Leuven, December 10th, 2009

Dear Editor,

Thank you for considering my paper with reference EHD-D-09-00126 titled " Decoupling between fundamental frequency and energy envelope of neonate cries ".

Although the decision letter indicates a major revision, the comments of the reviewer (reviewer 1) were more indicative of a minor revision. The email is shown below.

As mentioned by the reviewer the experience of the pain evaluator and physician for both MBPS assessment and venepuncture is important. As indicated in the revised tekst both activities were done by experienced staff. Also the fact that there is a substantial variation in the calculated variables of the acoustic emmission is highlighted. Furthermore, the book references have been checked and corrected as mentioned by the reviewer.

Hopefully these corrections are in compliance with the reviewers' vision.

Think you in advance,
Kind Regards,

Mitchell Silva

E-mail: Mitchell.silva@biw.kuleuven.be
Address: Catholic University of Leuven
Faculty of Bioscience Engineering
Biosystems Department
Measure, Model & Manage Bioresponses: M3-BIORES
Kasteelpark, Arenberg, B-3001
Heverlee / BELGIUM
Tel: ++32 (0)16 32 14 35
Fax: ++32 (0)16 32 14 80

MEASURE, MODEL & MANAGE BIORESPONSES (M3-BIORES)
KASTEELPARK ARGENBERG 30, B-3001 HEVERLEE

-----Original Message-----
From: ees.ehd.0.51b26.8abcb742@eesmail.elsevier.com
[mailto:ees.ehd.0.51b26.8abcb742@eesmail.elsevier.com] On Behalf Of Early Human Development
Sent: Tuesday, 17 November, 2009 23:46
To: Daniel Berckmans
Subject: Your Submission EHD-D-09-00126

Ms. Ref. No.: EHD-D-09-00126
Title: Decoupling between fundamental frequency and energy envelope of neonate cries
Early Human Development

Dear Prof. Daniel Berckmans,

Thank you for submitting your manuscript to Early Human Development. It has been carefully evaluated by our reviewers whom we consider to be experts in the field. The paper is not acceptable for publication in Early Human Development in its present form and will need a major revision and may require further evaluation by the reviewers. To assist you in revising your manuscript, we are enclosing the reviewer's comments. A final decision will be based on your response to the enclosed suggestions and criticisms.

When you submit a revised manuscript, please respond carefully to each of the comments by the reviewers and to editorial recommendations. You should include a detailed cover letter which delineates all changes to the manuscript and identifies the site in the manuscript where these changes have been made. The responses should be numbered to correspond to the reviewer's or editor's comments. You must clearly mark in the margin or highlight the places where the manuscript has been revised.

When submitting your revised manuscript, please ensure that you upload the source files (e.g. Word). Uploading a PDF file at this stage will create delays should your manuscript be finally accepted for publication. If your revised submission does not include the source files, we will contact you to request them.

To submit a revision, please go to http://ees.elsevier.com/ehd/ and login as an Author.
Your username is: Dberckmans
If you need to retrieve password details, please go to: http://ees.elsevier.com/EHD/automail_query.asp.

On your Main Menu page is a folder entitled "Submissions Needing Revision". You will find your submission record there.

We look forward to receiving your revised manuscript.

Sincerely,

Charles R. Rosenfeld, M.D.
Associate Editor
Early Human Development

Reviewers' comments:

Reviewer #1: Abstract: This paper is not only in accordance to the Journal rules but it is also clear and objective.

Introduction: General acknowledgement is provided and the purposes are clear.

IR. MITCHELL SILVA
TEL. +32 (0)16 32 14 35 FAX +32 (0)16 32 14 80
E-mail: mitchell.silva@biw.kuleuven.be
URL: http://www.m3-biores.be

MEASURE, MODEL & MANAGE BIORESPONSES (M3-BIORES)
KASTEELPARK ARGENBERG 30, B-3001 HEVERLEE

Methods: Were venepuncture and the MBPS Scale application done by the same evaluator? Although newborns pain scale are validated they depend on the evaluator's know-how, and the punction as well.

Results and discussion:
1. Tables and graphics are in accordance to the Journal rules.
2. According to published papers it is known that the stress, when facing pain, can make the newborn nervous system unstable. So, the acoustic emissions will be modified. Were there variations of acoustic characteristics during the evaluated period?

Conclusion: This paper shows a method for newborn pain cry evaluation and gives replies to research purpose.

References: At references 1, 4, 5 and 13 the sequence of date, place and press should be revised according to Vancouver style.

****

For further assistance, please visit our customer support site at http://epsupport.elsevier.com. Here you can search for solutions on a range of topics, find answers to frequently asked questions and learn more about EES via interactive tutorials. You will also find our 24/7 support contact details should you need any further assistance from one of our customer support representatives.

IR. MITCHELL SILVA
TEL. +32 (0)16 32 14 35 FAX +32 (0)16 32 14 80
E-mail: mitchell.silva@biw.kuleuven.be
URL: http://www.m3-biores.be

*Detailed Response to Reviewers

Answers to reviewers

Hereby the answers to the suggestions of the reviewer.

Reviewer #1:

Abstract: This paper is not only in accordance to the Journal rules but it is also clear and objective.

Introduction: General acknowledgement is provided and the purposes are clear.

Methods: Were venepuncture and the MBPS Scale application done by the same evaluator? Although newborns pain scale are validated they depend on the evaluator's know-how, and the punction as well.

Indeed, experience has an influence on both venepuncture and pain assessment. In the described study venepunctures were performed by a trained paediatrician, the MBPS was done by a dedicated evaluator trained for neonatal pain assessment based on the MBPS and was not involved in the venepuncture.

Page 6 line 3; Page 9 line7.

Results and discussion:

1. Tables and graphics are in accordance to the Journal rules.
2. **According to published papers it is known that the stress, when facing pain, can make the newborn nervous system unstable. So, the acoustic emissions will be modified. Were there variations of acoustic characteristics during the evaluated period?**

This is a good point. The variations in acoustic characteristics are indicated by the standard deviation of the calculated variables. The intra subject variablity of the correlation between fundamental frequency variation and energy contour shows a big standard deviation, which is related to a modified acoustic emission. The mean variation (standard deviation) of the fundamental frequency during the observation period is about 10% of the mean fundamental frequency (547 $\pm$52 Hz), while for the correlation coefficient this is on average 56% of the mean value (0.43$\pm$0.24). Although we did not measure the same individuals in various stress states to determine the change in variation of acoustic emission, the calculated standard deviations show there is a substantial variation in the acoustic features which are calculated during the observation of the infant.

Page 9 line 14; Page 10 line 4; Page 11 line 6.

Conclusion: This paper shows a method for newborn pain cry evaluation and gives replies to research purpose.

References: At references 1, 4, 5 and 13 the sequence of date, place and press should be revised according to Vancouver style.

The mentioned references have been adjusted:

1. Barr RG, Hopkins B, Green JA. Crying as a sign, a symptom and a signal., London: Mac Keith Press; 2000. pp. 13-14

4. Wolf P. The natural history of crying and other vocalization in early infancy. In: Foss B Determinants of infant behaviour. London: Methuen Press; 1969. pp. 81-109

5. Barr RG, Hopkins B, Green JA. Crying as a sign, a symptom and a signal. London: Mac Keith Press; 2000. pp. 143-146

13. Rangaraj M. Ramgayyan. Biomedical Signal Processing- A Case-Study approach. New York: John Wiley & Sons INC; 2002. pp. 203

Page 16 references 1,4,5 and 13.

Title:

Decoupling between fundamental frequency and energy envelope of neonate cries

Running title:

Decoupling in infant cries

List of authors:

Mitchell Silva[1], Bogdan Mijovic[4], Bea R.H. Van den Bergh[2], Karel Allegaert[3] , Jean-Marie Aerts[1], Sabine Van Huffel[4], Daniel Berckmans[1*]

Author Institutional Affiliations:

[1] Department of Biosystems, Katholieke Universiteit Leuven, Kasteelpark Arenberg 30, B-3001 Leuven, Belgium

[2] Department of Psychology, Tilburg University, P.O.Box 90153, 5000 LE Tilburg, The Netherlands and Department of Psychology, Katholieke Universiteit Leuven, Tiensestraat 102, B-3000 Leuven., Belgium

[3] Department of women and child, Group Biomedical Sciences, Katholieke Universiteit Leuven, Herestraat 49, B-3000 Leuven, Belgium

[4] Department of Electrical Engineering (ESAT), Kasteelpark Arenberg 10 bus 2446, B-3001 Leuven, Belgium

***Corresponding Author:**

Daniel Berckmans, Prof., PhD

Mail address:

Kasteelpark Arenberg 30, 3001 Leuven, Belgium

Telephone nr.: ++32 (0) 16 32 17 26

Fax: ++32 (0) 16 32 14 80

Email: Daniel.Berckmans@biw.kuleuven.be

*** 3. Ethical Statement**

Department of women and child
Group Biomedical Sciences
Katholieke Universiteit Leuven
Herestraat 49, B-3000 Leuven
Belgium

KATHOLIEKE
UNIVERSITEIT
LEUVEN

Leuven, July 9th, 2009

Ethical Statement

The study described in the paper titled "Decoupling between fundamental frequency and energy envelope of neonate cries" was approved by the Ethical Board of the University Hospitals (KULeuven, Belgium) and neonates were only included after written consent of the parents.

*Manuscript (without author names, details and affiliations)

Title:

Decoupling between fundamental frequency and energy envelope of neonate cries

Abstract:

Background: The presence of decoupling, i.e. the absence of coupling between fundamental frequency variation and energy envelope during phonetic crying, and its extent, reflects the degree of maturation of the central nervous system.

Aim: We hereby wanted to assess the existence and extent of decoupling in term neonates (neurodevelopmental relevance) and whether an association between decoupling and clinical pain expression could be unveiled (clinical relevance).

Study design: To assess decoupling in healthy term neonates during procedural pain, newborns were videotaped and crying was recorded during venous blood sampling. Besides acoustic analysis, pain expression was quantified based on the Modified Behavioral Pain Scale (MBPS).

Subjects: 47 healthy term neonates underwent venous blood puncture at the 3rd day of life.

Outcome measures: Beside the MBPS-score, the correlation coefficients were calculated between the fundamental frequency variation and energy envelope of the cries.

Results: Based on data collected in 47 healthy term neonates, correlation coefficients varied between 0.20 and 0.68. The degree of decoupling displayed extensive variability between the neonates and also in different cry bouts in a crying sequence within an individual neonate. A negative association was found between MBPS-value and

decoupling (r^2=-0.12), the same as for the intra-subject variability although less extensive (r^2=-0.02).

Conclusion: Decoupling only relates weakly with the amount of distress in 3 day old newborns, even though a great intra-subject variability is present. This study suggests that there is no evidence of extensive decoupling as the newborn still has to fully develop the control of larynx and abdominal muscles.

Key words:

cry analysis, decoupling, infant, maturation, Modified Behavioural Pain Scale

List of abbreviations:

AR	Auto-Regressive
DAN	Douleur Aigue du Nouveau-Né
F0	Fundamental Frequency
EE	Energy Envelope
MBPS	Modified Behavioural Pain Scale
RMS	Root Mean Square

Introduction

One of the most important tools for infants to communicate is vocalization. In neonates, vocalization is mainly crying. Crying is considered to be a gradual signal, variable in duration, in intensity and whose meaning needs to be interpreted in its context (1). The extent at which a newborn is stressed will also be reflected in its cry acoustics, causing different gradations in the cry signal, for example pitch or loudness. These gradations do not carry a specific message, but are more general, sudden and intense signals of stress.

From neurodevelopmental point of view, baby cries are the result of the interaction between control of different areas in the brain, respiratory control and vocal fold vibrations. At early stage, it is believed that a cry is the result of respiratory action and the effect of air going through a pipe, causing the vocal folds to vibrate, resulting in a cry bout (2), this is the vocalization produced during one expiration. The more the neural system matures, the more laryngeal control can be exerted resulting in manipulation and modulation of the cry signal, but observations on maturational aspects of the cry signal are limited and conflicting (3, 4). The development of the central neural system has been linked with the extent of vocalization control (4). Based on the cry production model of Golub described in the work of Moller and Schonweiler (2) and Barr *et al.* (5), decoupling between variations in fundamental frequency and energy envelope during crying indicates cortical control. Decoupling is referred to as the absence of coupling between fundamental frequency variation and energy envelope during phonetic crying. In this cry model, cry production is controlled by a three level processor structure, in which the lower level can be separated into the subglottal, glottal and supraglottal cry

production areas. The mid-processor is involved in physiological inputs like pain, blood levels and respiratory constraints, while the feed-back processes are part of the upper-processor level, especially auditory feedback.

Besides crying, the response in which a newborn reacts on an external impulse like pain can also be quantified with validated clinical scoring systems. Such clinical scoring systems are often multimodal and not only include scoring of the body movements, but also acoustic features of the cry signal (6, 7, 8, 9). Within a predefined context like pain, analysis of crying characteristics might therefore provide additional information (3,10).

Bellieni *et al.* were able to document that newborns with a higher DAN (Douleur Aigue du Nouveau-né) score during venous blood sampling (heel lancing) also had a higher fundamental frequency and a correlation was found between the pain intensity (DAN-score) and the normalized Root Mean Square value (RMS) (3). The observation that the normalized RMS value increases with the DAN score indicates that the stationary character of the overall cry intensity increases with increasing pain. This can be interpreted as the energy envelope showing little variations during a cry bout with increasing pain. Facchini *et al.* looked to the relation between the DAN score and the occurrence of noise patterns in the sound spectrograms (10). These noise patterns were chaotic and showed discontinuous parts in the crying signal, caused by a highly turbulent flow in the larynx. Results indicated that newborns with a higher DAN score showed more noise patterns.

To link cry characteristics with maturation, Wolf suggested that prior to one month of age infant cries are highly reflective and undifferentiated, but that shortly afterwards, cries progressively reflect various psycho-physiological states like hunger and pain (4). In

contrast, Bellieni *et al.* were able to document a correlation between cry features and DAN score in term neonates (3).

In the current study, we wanted to describe the dynamics within the baby cry and try to link these acoustic variables with clinical indicators of pain or stress expression in term neonates. We hereby wanted to investigate *(i)* whether decoupling can already be observed in term neonates (neurodevelopmental relevance), *(ii)* whether variability in decoupling could be quantified across the newborns and within the different cry bouts of one individual, and finally *(iii)* whether the degree of decoupling relates to clinical pain expression after venous blood sampling (clinical relevance).

Methods:

Clinical pain expression and vocalization were simultaneously recorded in term neonates in whom a venous blood sampling was performed on the 3rd day of life for routine metabolic screening. Venepunctures were performed by a trained paediatrician. All babies were born in the University Hospital of Leuven (UZLeuven), in the period of July 2005 until March 2006. The study was approved by the Ethical Board of the University Hospitals and neonates were only included after written consent of the parents. In order to participate in this project, the baby had to fulfil the following inclusion criteria: minimal gestational age at least 37 weeks, no complications during pregnancy, a 1- and 5- minutes APGAR score above 7, a minimal birth weight of 2.500 kg.

The vocalization produced by the newborn was recorded up to three minutes after the venous puncture, with a sampling frequency of 44.1 kHz at a distance of 0.5 m of the head of the baby. A cry from a baby consists of various cry bouts. Every cry bout is the result of the vibrations of the vocal folds during expiration. Due to the fact that the baby has to inspire, the cry bouts are separated by a pause. In infant cry analysis three types of vocalization are described (11): phonation (voiced cries), hyperphonation (high pitched cries) and disphonation (turbulence, voiceless cry). Hyperphonation is often characterized by a higher fundamental frequency, while in disphonation a clear fundamental frequency and its harmonics are absent. In order to evaluate the relation between the variation of fundamental frequency and energy envelope, only manually selected voiced cry bouts (phonation) are used with a clear fundamental frequency.

Examples of cry bouts from a recorded crying signal are shown in figure 1 and 2. Signal analysis was done with Matlab R14 (Mathworks) and Adobe Audition 2.0 (Adobe). For every cry signal, consisting of various cry bouts, the relation between fundamental frequency and energy envelope is investigated. First the signals were filtered using a 6th order Butterworth filter with lower cut-off frequencies 200 Hz and upper cut of frequency 2500 Hz, followed by a down sampling by a factor 8 of the signal to a sampling frequency of 5512 Hz. This allows us to investigate frequencies up to about 2700 Hz, under which the fundamental frequency is situated. The fundamental frequency was extracted by sliding a moving window (Hamming window of length 128 samples with 50% overlap) over the total cry bout. In every window, the fundamental frequency was derived, simultaneously with its energy envelope, using an autoregressive (AR) spectral estimation method (12). The order of the AR model will influence the amount of peaks that are to be modelled. The higher the order, the more estimated peaks. In our case, the model order was set to 20, based on the Akaike's Information Criterion (AIC). Figure 3 illustrates the estimated AR spectrum on top of the Power Spectral Density plot (spectrum) for a clearly phonated cry signal. Using peak detection, the fundamental frequency is extracted for every window. The energy envelope is derived from taking the absolute value of the total cry bout after which it is filtered by a low pass 3rd butterworth filter at 40 Hz (13). Afterwards the total energy envelope is windowed by the same window size as for the fundamental frequency extraction (128 samples) in which every window is represented by its mean to bring the energy envelope to the same sampling frequency as the fundamental frequency variation.

So, for every sliding window in a cry bout for which the AR spectrum is estimated, simultaneously the energy envelope was calculated. The result is shown in figure 1, in which the waveform is shown of a cry signal, together with its frequency variation and energy envelope (both normalized for visualization). In order to evaluate the degree of decoupling between the energy envelope and fundamental frequency (F0), correlation coefficient is calculated. The correlation coefficient *R(x,y)* between signals *x* and *y* is related to the covariance *C* in the following way:

$$R(x,y) = \frac{\mathrm{cov}(x,y)}{\sigma(x)\sigma(y)}$$

In this formulation cov(x,y) is the covariance between the vectors x and y, respectively the energy envelope and the fundamental frequency variation. The standard deviations of x and y are represented as σ(x) and σ(y). The higher the degree of coupling, the higher the correlation coefficient R will be. A coupled and a decoupled cry bout are provided in respectively figure 1 and 2. In the decoupled cry bout (fig. 2, R=0.12), there is an initial rise in fundamental frequency after which it decreases, while in the coupled example (fig. 1, R=0.77) the energy envelope and fundamental frequency tend to follow each other. Since this correlation coefficient between energy envelope and fundamental frequency quantifies decoupling, it decreases as decoupling increases (i.e. a correlation coefficient R = 0 means total decoupling while a correlation coefficient R = 1 means total coupling). As decoupling/coupling in a cry bout is characterized by a wide range of possible transitions between both states, it is more likely to speak about coupling (decoupling)

when the correlation coefficient goes towards 1 (0), instead of using a threshold value for the classification between the two classes.

Clinical pain expression was quantified based on the validated Modified Behavioural Pain Scale (MBPS) (14,15). This is an easy applicable pain scale, based on visual and auditive scoring of facial expression, crying and body movements, each with a specific gradation. Summing the different sub scores results in a total score between 0 and 10 (table 1). The MBPS was done by a dedicated evaluator trained for neonatal pain assessment based on the MBPS and was not involved in the venepuncture.

Results

47 healthy term neonates (20 boys, 27 girls), with a mean weight of 3.3±0.4 kg were included, resulting in the recording of 1330 cry bouts. After selecting only those cry bouts with a clear phonation to enable the quantification of fundamental frequency variation, 1175 cry bouts remain. Mean MBPS was 7, and varied between 2 and 10. The mean number of cry bouts recorded in the 3 minutes time interval was 28 (range 4 – 97). The mean gestational age was 39 weeks and 4 days (±1 week and 2 days). Mean fundamental frequency was 547 (±52) Hz. The highest fundamental frequency was observed at 843 Hz, the lowest at 200 Hz.

Neurodevelopmental outcome: The degree of decoupling between fundamental frequency and energy envelope during a cry showed extensive inter-individual

variability. In this cohort, the average correlation coefficients per subject calculated over all the cry bouts of the corresponding baby, recorded during the 3 minute test, varied between 0.20 and 0.68. A great variability between the degree of coupling/decoupling over all neonates has been observed, with a mean correlation coefficient of 0.43±0.24.

When looking more in detail to the correlations of the cry bouts in a crying sequence, we also observed extensive intra-individual variability, expressed as the standard deviation over all cry bouts from one baby. This is illustrated in figure 5, where the correlation coefficients are shown for the 34 cry bouts recorded in one neonate. Although this baby displays a strong correlation between fundamental frequency and energy envelope up to 0.69, some cry bouts only have a low correlation coefficient(i.e. only 0.03) indicating higher degree of decoupling. In this case the mean correlation coefficient over all the 34 cry bouts is 0.29±0.19. Over all the subjects, the standard deviation fluctuates between 0.069 and 0.342, indicating great intra-subject variability.

Clinical outcome: When evaluating the number of cries against the MBPS-score, a weak and statistically not significant positive correlation was observed (r^2=0.01). It is not significant as the 95% confidence interval of the slope of the regression line, called β, holds 0 within its range, being -0.7±113. Nor mean fundamental frequency nor standard deviation of the fundamental frequency during a crying sequence shows a relation with MBPS score (respectively r^2=-0,01 and r^2=0,03). When plotting the MBPS values against the correlation coefficients (figure 5), no strong relation could be found, although significant (β= -0.0144±0.0018). The linear regression line has an r^2 of -0.12. The MBPS shows a weak, non significant relation with the individual standard deviations (r^2=-0.02).

Discussion and conclusion

Based on crying characteristics collected during 47 procedures in term neonates, we were able to quantify decoupling and illustrate that it has an important inter- and intra-individual variability and that the degree of decoupling has a slight negative correlation with MBPS (r^2=-0.12), while the intra-subject variability (defined as the standard deviation of the correlation coefficient quantifying the degree of decoupling) shows a less extensive negative relation with increasing MBPS value (r^2=0.02).

Neurodevelopmental aspects: The phenomenon of decoupling can be explained by the fact that the fundamental frequency and the amplitude envelope (energy envelope) are controlled by different muscles and nerves (16). The fundamental frequency is controlled by airflow and the cricothyroid muscle, innervated by the external branch of the superior laryngeal nerve. This is a branch of the nervus vagus of which the motorneurons are located in the rostral part of the nucleus ambiguous in the brain stem. In contrast, the amplitude envelope mainly relates to abdominal wall muscles, which is innervated by the nerves T2-L3. The internal and external oblique muscles are activated before the mm. transverses and rectus abdominis. While the oblique muscles and the m. transverses are well correlated in their activity with the amplitude envelope of phonation, there is only a loose relation between m. rectus activity and amplitude envelope (17).

Kent reports that the early divergence between speech and non-speech motor activities indicates that separate neural control systems are established in infancy (18). Wermke *et al.* (19) documented a progressive increase in the ability of decoupling during crying in

twins at age 8 -9 weeks, 15-17 weeks and 23 -24 weeks (postnatal age). Although the increase of the degree of decoupling has been observed, it was not clear to which extent three day old term neonates were able to decouple their energy envelope from their fundamental frequency. The results presented in this paper have quantified the extent of this decoupling. In total, the newborns showed an average correlation coefficient of 0.43, and the standard deviations of the correlation coefficients were up to 76% of their mean value, indicating high inter-and intra-subject variability. This suggests that there is no evidence of extensive controlled decoupling, probably as the newborn still has to fully develop the control of larynx and abdominal muscles. As mentioned earlier, the air expired from the lungs causes the vocal folds to vibrate, producing a fundamental frequency. According to the neural cry model discussed by Porter *et al.* (20), the modulation of this fundamental frequency is related to the dual innervation of the larynx by both parasympathetic and sympathetic input of the autonomic nervous system. Normal neurological development is characterized by a balance between these excitatory (sympathetic) and inhibitory (parasympathetic) forces. This whole process makes it unclear whether decoupling/coupling is clearly to be separated into cortical and subcortical processes, which might explain the high variability in the results.

Clinical aspects: When comparing the extent of clinical pain expression (MBPS score) with the degree of decoupling, a slight negative relation could be found. This is shown in figure 4 where a negative trend can be observed, although with low correlation (r^2=0.12). This might indicate that the degree of decoupling (and hence developmental status of the laryngeal control ability) in three days old newborns does not hold in great extent

information about the stress reactivity expressed in MBPS values. In the past, other cry characteristics like fundamental frequency or oscillations have been linked with pain expression (3, 10). Bellieni *et al.* documented that the cry characteristics in neonates with a DAN score higher than 8 (higher reactivity) had a higher fundamental frequency (3). The reason why this is not observed in our experiment is that Bellieni et al. used the heel-prick to evoke crying, a more severe blood sampling technique which leads to a higher excitation of the infant compared with our method. Fundamental frequency is also related to the listener's perception of a cry bout. Protopapas and Eimas (21) have shown that fundamental frequency and jitter are related to the listeners' perception, but results were inconclusive for the rise time (the time from the offset of phonation until the pitch reaches its maximum value). As the rise time is related to the variation of fundamental frequency in a cry bout, the conclusions might be similar for the degree of decoupling, but this is yet to be investigated as the quantification of decoupling also involves the variation in energy envelope.

In this study, it also has been shown that the intra-subject variability –expressed as the standard deviation of the degree of decoupling - shows a very weak negative relation with MBPS value (r^2=0.02). One might interpret the standard deviation within a subject with the consistency by which a cry bout it produced. As the ratio of the standard deviation of the correlation coefficients and the mean correlation coefficient is large in the studied subjects, it can be said that the cry bouts are produced with a low consistency.

Facchini *et al.* observed increased occurrence of noise patterns with high DAN scores, more specific with DAN scores between 8 and 10 (10). These authors hereby postulated

that noise patterns could be seen as distortions in the oscillating system caused by large variations in the vocal fold tension and tissue constants in distressed babies. This would imply that an increase in pain induces a chaotic transition in the vocal fold oscillation leading to noise patterns. As disphonation does not show a clear fundamental frequency, the AR method used in this study will not give a clear frequency contour. This is why the AR method for F0 estimation has been applied for cry phonations with clear fundamental frequency variations, and not for disphonation. The absolute values of the fundamental frequencies, obtained in this study via AR spectra, are in the same range as described in literature (3, 10).

In conclusion, the method described in this paper is able to quantify decoupling and showed extensive inter- and intra-subject variability. Only a weak negative relation was found between decoupling and clinical reactivity (MBPS-values) during procedural pain, as well as for the relation between the inter-subject variability and MBPS values, but less extensive. The results show that decoupling is not fully present in term neonates, and that the intensity of pain expression is to some extent related to the degree of decoupling. The consistency in which the cry bouts are produced shows a weak negative relation with the clinical reactivity. We conclude that decoupling is not implicitly linked to pain, suggesting that the method described in this paper might be relevant for neurodevelopmental rather than clinical assessment during venous blood sampling. For future research, it would be interesting to further describe the evolution in cry complexity and its covariates during the first few months of development, as the laryngeal muscles are still to be further developed in the following months after birth.

Acknowledgements

The authors wish to thank all the families for participating in the study and Sofie De Vos, Marijke Fraussen, Elke Maes, May Magnus, Sylvia Pinna Puissant, Annelies Simons and Sara Weckhuyzen for gathering the data. The clinical research of K Allegaert is supported by the Fund for Scientific Research, Flanders (Belgium) (FWO Vlaanderen) by a Fundamental Clinical Investigatorship (1800209N). B. Mijovic and S. Van Huffel are supported by the Research Council KUL: GOA AMBioRICS, CoE EF/05/006, IDO 05/010 and the Belgian Federal Science Policy Office: IUAP P6/04.

References

1. Barr RG, Hopkins B, Green JA. Crying as a sign, a symptom and a signal., London: Mac Keith Press; 2000. pp. 13-14

2. Moller S, Schonweiler R. Analysis of infant cries for the early detection of hearing impairment. Speech Commun 1999; 28:175-193

3. Bellieni CV, Sisto R, Codelli BM, Buonocore G. Cry features reflect pain intensity in term newborns: an alarm threshold. Pediatr Res 2004; 55: 142-146

4.Wolf P. The natural history of crying and other vocalization in early infancy. In: Foss B Determinants of infant behaviour. London: Methuen Press; 1969. pp. 81-109

5. Barr RG, Hopkins B, Green JA. Crying as a sign, a symptom and a signal. London: Mac Keith Press; 2000. pp. 143-146

6 Braungart-Rieker J, Stifter C. Infants' Responses to Frustrating Situations: Continuity and Change in Reactivity and Regulation. Child Dev 1996; 67: 1767-1779

7. Hummel P, van Dijk M. Pain assessment: Current status and challenges. Semin Fetal Neonatal Med 2006; 11:237-245

8. Holsti L, Grunau RE. Initial validation of the Behavioral Indicators of Infant Pain (BIIP). Pain 2007; 132:264-272

9. Lehr VT, Zeskind PS, Ofenstein JP, Cepeda E, Warrier I, Aranda JV. Neonatal facial coding system scores and spectral characteristics of infant crying during newborn circumcision. Clin J Pain 2007; 23:417-424

10. Fecchini A, Bellieni CV, Marchettini N, Pulselli FM, Tiezzi EB. Relating pain intensity of newborns to onset of nonlinear phenomena in cry recordings. Phys Lett A 2005; 338: 332-337

11. LaGasse L, Neal A, Lester B. Assessment of infant cry: acoustic cry analysis and parental perception. Ment Retard Dev Disabil Res Rev 2005; 11: 83-93

12 Pardey J, Roberts S and Tarassenko L. A review of parametric modelling techniques for EEG analysis. Medical Engineering & Physics 1996; 18: 2-11

13. Rangaraj M. Ramgayyan. Biomedical Signal Processing- A Case-Study approach. New York: John Wiley & Sons INC; 2002. pp. 203

14. Ipp M, Taddio A, Sam J, Gladbach M, Parkin PC. Vaccine-related pain: randomised controlled trial of two injection techniques. Arch Dis Child 2007; 92:1105-1108

15. Taddio A, Nulman I, Goldbach M, Ipp M, Koren G. Use of lidocaine-prilocaine cream for vaccination pain in infants. J Pediatr 1994; 124: 643-648

16. Yoshida Y, Tanaka Y, Saito T, Shimazaki T, Hirano M. Peripheral nervous system in the larynx. An anatomical study of the motor, sensory and autonomic nerve fibers. Folia Phoniatr (Basel) 1992; 44:194–219

17. Jürgens U. Neural pathways underlying vocal control. Neurosci Biobehav Rev 2002; 26:235-258

18. Kent RD. Research on speech motor control and its disorders - A review and prospective. J Commun Disord 2000; 33:391-427

19. Wermke K, Mende W, Manfredi C, Bruscaglioni P. Developmental aspects of infant's cry melody and formants. Med Eng Phys 2002; 24:501-514

20. Porter FL, Porges SW, Marchall RE. Newborn pain cries and vagal tone-parallel changes in response to circumcision. Child Dev 1988; 59: 495-505

21. Protopapas A, Eimas PD. Perceptual differences in infant cries revealed by modifications of acoustic features. J Acoust Soc Am 1997; 102: 3723-3734

Figure legends:

Figure 1: coupled cry bout showing its waveform with corresponding energy envelope and fundamental frequency variation (both normalized for visualization). The cross correlation coefficient R is 0.77.

Figure 2: uncoupled cry bout showing its waveform with corresponding energy envelope and fundamental frequency variation (both normalized for visualization). The cross correlation coefficient R is 0.12.

Figure 3: AR model based and corresponding PSD spectrum from a phonated cry signal: the first peak is the fundamental frequency, the following peaks the harmonics.

Figure 4: MBPS values versus the mean correlation coefficient of all cry bouts for all the subjects.

Figure 5: intra-subject variability is shown by the variation in correlation coefficient between different cry bouts of one crying signal. In this case the correlation varied between 0.03 and 0.70.

Table Legends:

Table 1: the three scoring components of the MBPS are facial expression, cry and movements. The maximum possible MBPS value is 10.

Table 2: data overview.

6. Table(s)

Table 1: the three scoring components of the MBPS are facial expression, cry and movements. The maximum possible MBPS value is 10.

Facial expression	score
Definite positive expression	0
Neural expression	1
Slightly negative expression (i.e. grimace)	2
Definite negative expression (i.e. furrowed brows, eyes close tightly)	3
Cry	
Laughing or giggling	0
Not crying	1
Moaning, quiet vocalizing or gentle or whimpering cry	2
Full-lunged cry or sobbing	3
Full-lunged cry, clearly more than baseline	4
Movements	
Usual movements and activity	0
Resting and relaxed	1
Partial movement or attempt to avoid pain by withdrawing	2
Agitation with complex movements or rigidity	3

Table 2: data overview

subject	voiced cry	total cry	MBPS	mean corr coeff ± st. dev.	mean FREQ ± st. dev.
1	39	66	8	0.494 ± 0.293	481 ± 33
2	63	71	9	0.277 ± 0.229	461 ± 40
3	28	30	3	0.480 ± 0.256	557 ± 28
4	37	60	5	0.515 ± 0.262	548 ± 28
5	7	8	4	0.468 ± 0.200	509 ± 28
6	58	64	7	0.394 ± 0.255	578 ± 31
7	27	33	10	0.472 ± 0.207	458 ± 26
8	18	19	9	0.355 ± 0.259	462 ± 27
9	6	6	3	0.680 ± 0.271	533 ± 26
10	41	42	10	0.406 ± 0.237	455 ± 37
11	12	13	8	0.306 ± 0.195	503 ± 29
12	33	34	10	0.381 ± 0.247	432 ± 194
13	19	21	8	0.311 ± 0.217	639 ± 24
14	36	48	10	0.430 ± 0.266	480 ± 26
15	34	37	6	0.290 ± 0.197	537 ± 25
16	14	15	3	0.486 ± 0.245	495 ± 25
17	11	11	9	0.448 ± 0.279	537 ± 36
18	4	4	8	0.202 ± 0.069	472 ± 35
19	8	12	8	0.540 ± 0.206	707 ± 32
20	19	23	6	0.337 ± 0.268	462 ± 32
21	16	26	8	0.565 ± 0.258	511 ± 22
22	11	11	8	0.377 ± 0.271	467 ± 25
23	25	29	10	0.369 ± 0.248	472 ± 27
24	11	12	9	0.330 ± 0.230	843 ± 47
25	36	40	10	0.548 ± 0.269	474 ± 26
26	8	8	4	0.470 ± 0.192	463 ± 31
27	30	31	8	0.424 ± 0.237	525 ± 44
28	15	17	4	0.377 ± 0.222	478 ± 21
29	28	29	8	0.508 ± 0.277	480 ± 22
30	8	8	7	0.434 ± 0.342	467 ± 31
31	70	74	8	0.363 ± 0.249	200 ± 79
32	91	97	7	0.422 ± 0.247	521 ± 27
33	33	36	10	0.270 ± 0.187	531 ± 70
34	37	37	3	0.596 ± 0.224	601 ± 41
35	8	8	7	0.319 ± 0.206	739 ± 36
36	22	24	7	0.451 ± 0.248	561 ± 27
37	21	22	5	0.601 ± 0.294	663 ± 42
38	9	10	8	0.420 ± 0.306	663 ± 58
39	13	13	10	0.485 ± 0.260	610 ± 28
40	10	12	8	0.517 ± 0.307	661 ± 80
41	16	17	4	0.469 ± 0.294	608 ± 32
42	27	28	8	0.552 ± 0.255	621 ± 76
43	37	37	8	0.444 ± 0.279	673 ± 32
44	29	30	10	0.391 ± 0.253	679 ± 37
45	30	34	10	0.389 ± 0.224	593 ± 32
46	6	6	2	0.445 ± 0.342	709 ± 36
47	14	17	8	0.583 ± 0.222	583 ± 29

figure1
Click here to download high resolution image




figure2
Click here to download high resolution image




figure3
Click here to download high resolution image




figure4
Click here to download high resolution image




figure5
Click here to download high resolution image

