NEW UNIQUENESS CONDITIONS FOR THE CANONICAL POLYADIC DECOMPOSITION OF THIRD-ORDER TENSORS

MIKAEL SØRENSEN * † AND LIEVEN DE LATHAUWER * †

Abstract. The uniqueness properties of the Canonical Polyadic Decomposition (CPD) of higher-order tensors make it an attractive tool for signal separation. However, CPD uniqueness is not yet fully understood. In this paper, we first present a new uniqueness condition for a CPD where one of the factor matrices is assumed known. We also show that this result can be used to obtain a new overall uniqueness condition for the CPD. In signal processing the CPD factor matrices are often constrained. Building on the preceding results, we provide a new uniqueness condition for a CPD with a columnwise orthonormal factor matrix, representing uncorrelated signals. We also obtain a new uniqueness condition for a CPD with a partial Hermitian symmetry, useful for tensors in which covariance matrices are stacked, which are common in statistical signal processing. We explain that such constraints can lead to more relaxed uniqueness conditions. Finally, we provide an inexpensive algorithm for computing a CPD with a known factor matrix that is also useful for the computation of the full CPD.

Key words. tensor, polyadic decomposition, parallel factor (PARAFAC), canonical decomposition (CANDECOMP), canonical polyadic decomposition.

1. Introduction. Tensor decompositions are finding more and more applications in signal processing, data analysis and machine learning. For instance, Canonical Polyadic Decomposition (CPD) is becoming a basic tool for signal separation. Essentially, this is due to the fact that CPD is unique under mild conditions, compared to decomposition of a matrix in rank-1 terms. Thanks to their uniqueness, the rank-1 terms are easily associated with interpretable data components. Numerous applications have been reported in independent component analysis, exploratory data analysis, wireless communication, radar, chemometrics, psychometrics, sensor array processing, and so on [1, 2, 7, 19, 20, 16, 18, 13, 14]. However, the understanding of uniqueness is lagging behind the use of CPD in practice.

Many uniqueness conditions for the CPD such as those developed in [15, 12, 26, 8, 9] are based on Kruskal's permutation lemma [15]. To prove uniqueness of a CPD, one may first prove the uniqueness of one factor matrix. In the next step, overall CPD uniqueness is obtained from the uniqueness of a CPD with a known factor matrix. The former problem has been thoroughly studied in [8]. In this paper we focus on the latter problem.

Besides uniqueness, the problem of studying CPD with a known factor matrix has direct relevance in applications. As an example, we mention that several wireless communication systems based on tensor-coding have been proposed (e.g. [3, 17, 4, 5]) since their conception in [21]. It suffices to say that many tensor-based wireless communication systems essentially rely on computing a CPD with a known factor. However, mainly optimization-based methods such as Alternating Least Squares (ALS) have

*Group Science, Engineering and Technology, KU Leuven - Kulak, E. Sabbelaan 53, 8500 Kortrijk, Belgium, KU Leuven - E.E. Dept. (ESAT) - STADIUS Center for Dynamical Systems, Signal Processing and Data Analytics, and iMinds Medical IT Department, Kasteelpark Arenberg 10, B-3001 Leuven-Heverlee, Belgium. {Mikael.Sorensen, Lieven.DeLathauwer}@kuleuven-kulak.be.

†Research supported by: (1) Research Council KU Leuven: GOA-MaNet, CoE EF/05/006 Optimization in Engineering (OPTEC), CIF1, STRT1/08/23, (2) F.W.O.: project G.0427.10N, G.0830.14N, G.0881.14N, (3) the Belgian Federal Science Policy Office: IUAP P7 (DYSCO II, Dynamical systems, control and optimization, 2012-2017), (4) EU: ERC advanced grant no. 339804 (BIOTENSORS).

been considered. Such iterative methods are not guaranteed to find the decomposition, even in the exact case. Thus, an optimization-based method can suffer from slow convergence (many iterations may be needed) and local minima (many initializations may be needed). On the other hand, an algebraic method is guaranteed to find the decomposition in the exact case. For sufficiently high Signal-to Noise Ratio (SNR), algebraic methods can provide an inexpensive but accurate estimate of the solution that can be used to initialize an optimization-based algorithm. Hence, the development of an algebraic method for computing the CPD with a known factor matrix is another relevant problem that will be addressed in this paper. For this reason we will develop a constructive uniqueness proof for the CPD with a known factor matrix.

The contributions of this paper are the following. We first present a new constructive uniqueness condition for a CPD with a known factor matrix that leads to more relaxed conditions than those obtained in [9] and is eligible in cases where none of the involved factors have full column rank. Based on this result we propose a new relatively easy-to-check deterministic overall uniqueness condition for the CPD that is comparable with recent relaxed deterministic overall CPD uniqueness conditions [9]. In the supplementary material we present a partial uniqueness variant. In tensor-based signal processing CPD is often constrained. Uncorrelated signals may translate into a columnwise orthonormal factor matrix [23] while stacking covariance matrices may lead to a CPD with a partial Hermitian symmetry (e.g. [2, 7]). We provide a new uniqueness condition for a CPD with a columnwise orthonormal factor matrix. We also provide a new uniqueness condition for a CPD with a partial Hermitian symmetry. We explain that such constrained CPDs can be unique under milder conditions than their unconstrained counterparts. Finally, we present an algorithm for computing a CPD with a known factor matrix that does not cost much more than a single ALS iteration. Numerical experiments demonstrate that this inexpensive algorithm provides a good initialization for a subsequent optimization-based method such as ALS in difficult cases with modest to high SNR.

The paper is organized as follows. The rest of the introduction presents our notation. Sections 2 and 3 briefly review the CPD and the CPD with a known factor matrix, respectively. Section 4 presents a new uniqueness condition and an algorithm for a CPD with a known factor matrix. Next, in section 5 we present new uniqueness conditions for the overall CPD and for some variants. Numerical experiments are reported in section 6. Section 7 concludes the paper. We also mention that in the supplementary material a new partial uniqueness condition for CPD and an efficient implementation of the ALS method for CPD with a known factor matrix are reported.

1.1. Notation. Vectors, matrices and tensors are denoted by lower case boldface, upper case boldface and upper case calligraphic letters, respectively. The rth column vector of $\mathbf{A}$ is denoted by $\mathbf{a}_r$. The symbols $\otimes$ and $\odot$ denote the Kronecker and Khatri-Rao products, defined as

$$\mathbf{A} \otimes \mathbf{B} \triangleq \begin{bmatrix} a_{11}\mathbf{B} & a_{12}\mathbf{B} & \dots \\ a_{21}\mathbf{B} & a_{22}\mathbf{B} & \dots \\ \vdots & \vdots & \ddots \end{bmatrix}, \qquad \mathbf{A} \odot \mathbf{B} \triangleq \begin{bmatrix} \mathbf{a}_1 \otimes \mathbf{b}_1 & \mathbf{a}_2 \otimes \mathbf{b}_2 & \dots \end{bmatrix},$$

in which $(\mathbf{A})_{mn} = a_{mn}$. The Hadamard product is denoted by $*$ and satisfies $(\mathbf{A} * \mathbf{B})_{mn} = a_{mn}b_{mn}$. The outer product of N vectors $\mathbf{a}^{(n)} \in \mathbb{C}^{I_n}$ is denoted by $\mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \cdots \circ \mathbf{a}^{(N)} \in \mathbb{C}^{I_1 \times I_2 \times \cdots \times I_N}$, such that

$$\left(\mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \cdots \circ \mathbf{a}^{(N)}\right)_{i_1,i_2,\dots,i_N} = a^{(1)}_{i_1} a^{(2)}_{i_2} \cdots a^{(N)}_{i_N}.$$

The identity matrix, all-zero matrix and all-zero vector are denoted by $\mathbf{I}_M \in \mathbb{C}^{M\times M}$, $\mathbf{0}_{M,N} \in \mathbb{C}^{M\times N}$ and $\mathbf{0}_M \in \mathbb{C}^M$, respectively. The symbol δ_{ij} denotes the Kronecker delta function, equal to 1 if $i = j$ and 0 if $i \neq j$.

The real part, imaginary part, transpose, conjugate, conjugate-transpose, inverse, Moore-Penrose pseudo-inverse, Frobenius norm, determinant, range and kernel of a matrix are denoted by $\mathrm{Re}\{\cdot\}$, $\mathrm{Im}\{\cdot\}$, $(\cdot)^T$, $(\cdot)^*$, $(\cdot)^H$, $(\cdot)^{-1}$, $(\cdot)^\dagger$, $\|\cdot\|_F$, $|\cdot|$, $\mathrm{range}(\cdot)$ and $\ker(\cdot)$, respectively.

The rank of a matrix $\mathbf{A}$ is denoted by $r(\mathbf{A})$ or $r_{\mathbf{A}}$. The k-rank of a matrix $\mathbf{A}$ is denoted by $k(\mathbf{A})$ or $k_{\mathbf{A}}$. It is equal to the largest integer $k_{\mathbf{A}}$ such that every subset of $k_{\mathbf{A}}$ columns of $\mathbf{A}$ is linearly independent. The cardinality of a set S is denoted by $\mathrm{card}(S)$.

Given $\mathbf{A} \in \mathbb{C}^{I\times J}$, then $\mathrm{Vec}(\mathbf{A}) \in \mathbb{C}^{IJ}$ will denote the column vector defined by $(\mathrm{Vec}(\mathbf{A}))_{i+(j-1)I} = (\mathbf{A})_{ij}$. Given $\mathbf{a} \in \mathbb{C}^{IJ}$, then the reverse operation is $\mathrm{Unvec}(\mathbf{a}) = \mathbf{A} \in \mathbb{C}^{I\times J}$ such that $(\mathbf{a})_{i+(j-1)I} = (\mathbf{A})_{ij}$. $D_k(\mathbf{A}) \in \mathbb{C}^{J\times J}$ denotes the diagonal matrix holding row k of $\mathbf{A} \in \mathbb{C}^{I\times J}$ on its diagonal.

Matlab index notation will be used for submatrices of a given matrix. For example, $\mathbf{A}(1:k,:)$ represents the submatrix of $\mathbf{A}$ consisting of the rows from 1 to k of $\mathbf{A}$. Let $\mathbf{T} \in \mathbb{C}^{I\times R}$, then $\underline{\mathbf{T}} = \mathbf{T}(1:I-1,:) \in \mathbb{C}^{(I-1)\times R}$, i.e., $\underline{\mathbf{T}}$ is obtained by deleting the bottom row of $\mathbf{T}$.

The matrix that orthogonally projects on the orthogonal complement of the column space of $\mathbf{A} \in \mathbb{C}^{I\times J}$ is denoted by

$$\mathbf{P}_{\mathbf{A}} = \mathbf{I}_I - \mathbf{F}\mathbf{F}^H \in \mathbb{C}^{I\times I},$$

where the column vectors of $\mathbf{F}$ constitute an orthonormal basis for $\mathrm{range}(\mathbf{A})$.

Let $C_n^k = \frac{n!}{k!(n-k)!}$ denote the binomial coefficient. The k-th compound matrix of $\mathbf{A} \in \mathbb{C}^{m\times n}$ is denoted by $C_k(\mathbf{A}) \in \mathbb{C}^{C_m^k \times C_n^k}$ and its entries correspond to the k-by-k minors of $\mathbf{A}$ ordered lexicographically. As an example, let $\mathbf{A} \in \mathbb{C}^{4\times 3}$, then

$$C_2(\mathbf{A}) = \begin{bmatrix} |\mathbf{A}([1,2],[1,2])| & |\mathbf{A}([1,2],[1,3])| & |\mathbf{A}([1,2],[2,3])| \\ |\mathbf{A}([1,3],[1,2])| & |\mathbf{A}([1,3],[1,3])| & |\mathbf{A}([1,3],[2,3])| \\ |\mathbf{A}([1,4],[1,2])| & |\mathbf{A}([1,4],[1,3])| & |\mathbf{A}([1,4],[2,3])| \\ |\mathbf{A}([2,3],[1,2])| & |\mathbf{A}([2,3],[1,3])| & |\mathbf{A}([2,3],[2,3])| \\ |\mathbf{A}([2,4],[1,2])| & |\mathbf{A}([2,4],[1,3])| & |\mathbf{A}([2,4],[2,3])| \\ |\mathbf{A}([3,4],[1,2])| & |\mathbf{A}([3,4],[1,3])| & |\mathbf{A}([3,4],[2,3])| \end{bmatrix}.$$

See [10, 8] for a discussion of compound matrices.

2. Canonical Polyadic Decomposition (CPD). Consider the third-order tensor $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$. We say that $\mathcal{X}$ is a rank-1 tensor if it is equal to the outer product of some non-zero vectors $\mathbf{a} \in \mathbb{C}^I$, $\mathbf{b} \in \mathbb{C}^J$ and $\mathbf{c} \in \mathbb{C}^K$ such that $x_{ijk} = a_i b_j c_k$. Decompositions into rank-1 terms are called polyadic decompositions:

$$\mathcal{X} = \sum_{r=1}^{R} \mathbf{a}_r \circ \mathbf{b}_r \circ \mathbf{c}_r \,. \tag{2.1}$$

The rank of a tensor $\mathcal{X}$ is equal to the minimal number of rank-1 tensors that yield $\mathcal{X}$ in a linear combination. Assume that the rank of $\mathcal{X}$ is R, then (2.1) is called the CPD of $\mathcal{X}$. Let us stack the vectors $\{\mathbf{a}_r\}$, $\{\mathbf{b}_r\}$ and $\{\mathbf{c}_r\}$ into the matrices

$$\mathbf{A} = [\mathbf{a}_1, \ldots, \mathbf{a}_R] \in \mathbb{C}^{I\times R}, \quad \mathbf{B} = [\mathbf{b}_1, \ldots, \mathbf{b}_R] \in \mathbb{C}^{J\times R}, \quad \mathbf{C} = [\mathbf{c}_1, \ldots, \mathbf{c}_R] \in \mathbb{C}^{K\times R}.$$

The matrices $\mathbf{A}$, $\mathbf{B}$ and $\mathbf{C}$ are called CPD factor matrices. Let $\mathbf{X}^{(i\cdot\cdot)} \in \mathbb{C}^{J\times K}$ denote the matrix such that $\left(\mathbf{X}^{(i\cdot\cdot)}\right)_{jk} = x_{ijk}$, then $\mathbf{X}^{(i\cdot\cdot)} = \mathbf{B}D_i\left(\mathbf{A}\right)\mathbf{C}^T$ and

$$\mathbb{C}^{IJ\times K} \ni \mathbf{X}_{(1)} \triangleq \left[\mathbf{X}^{(1\cdot\cdot)T}, \ldots, \mathbf{X}^{(I\cdot\cdot)T}\right]^T = \left(\mathbf{A} \odot \mathbf{B}\right)\mathbf{C}^T. \tag{2.2}$$

Similarly, let the matrices $\mathbf{X}^{(\cdot j\cdot)} \in \mathbb{C}^{K\times I}$ be constructed such that $\left(\mathbf{X}^{(\cdot j\cdot)}\right)_{ki} = x_{ijk}$, then $\mathbf{X}^{(\cdot j\cdot)} = \mathbf{C}D_j\left(\mathbf{B}\right)\mathbf{A}^T$ and

$$\mathbb{C}^{JK\times I} \ni \mathbf{X}_{(2)} \triangleq \left[\mathbf{X}^{(\cdot 1\cdot)T}, \ldots, \mathbf{X}^{(\cdot J\cdot)T}\right]^T = \left(\mathbf{B} \odot \mathbf{C}\right)\mathbf{A}^T. \tag{2.3}$$

Finally, let $\mathbf{X}^{(\cdot\cdot k)} \in \mathbb{C}^{I\times J}$ satisfy $\left(\mathbf{X}^{(\cdot\cdot k)}\right)_{ij} = x_{ijk}$, then $\mathbf{X}^{(\cdot\cdot k)} = \mathbf{A}D_k\left(\mathbf{C}\right)\mathbf{B}^T$ and

$$\mathbb{C}^{IK\times J} \ni \mathbf{X}_{(3)} \triangleq \left[\mathbf{X}^{(\cdot\cdot 1)T}, \ldots, \mathbf{X}^{(\cdot\cdot K)T}\right]^T = \left(\mathbf{C} \odot \mathbf{A}\right)\mathbf{B}^T. \tag{2.4}$$

2.1. Uniqueness conditions for one factor matrix of a CPD. A factor matrix, say $\mathbf{C}$, of the CPD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ is said to be unique if it can be determined up to the inherent column scaling and permutation ambiguities from $\mathcal{X}$. More formally, the factor matrix $\mathbf{C}$ is unique if alternative factor matrices $\widehat{\mathbf{C}}$ satisfy the condition

$$\widehat{\mathbf{C}} = \mathbf{C}\mathbf{P}\Delta_{\mathbf{C}}\,,$$

where $\mathbf{P}$ is a permutation matrix and $\Delta_{\mathbf{C}}$ is a nonsingular diagonal matrix. Based on Kruskal's permutation lemma [15] and the properties of compound matrices [10, 8], the following Theorem 2.1 guarantees the uniqueness of one factor matrix of a CPD. Theorem 2.1 is one of the most relaxed and yet quite easy-to-check sufficient (but not necessary) uniqueness conditions that have been reported in the literature. Consequently, we will use Theorem 2.1 to obtain new overall uniqueness results.

THEOREM 2.1. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in* (2.1). *If*

$$\begin{cases} k_{\boldsymbol{C}} \geq 1, \\ \min\left(I, J\right) \geq R - r_{\boldsymbol{C}} + 2, \\ C_{R-r_{\boldsymbol{C}}+2}\left(\boldsymbol{A}\right) \odot C_{R-r_{\boldsymbol{C}}+2}\left(\boldsymbol{B}\right) \;\; \textit{has full column rank,} \end{cases} \tag{2.5}$$

then the rank of $\mathcal{X}$ is R and the factor matrix $\boldsymbol{C}$ is unique [8].

We refer to [8] and references therein for other conditions that guarantee the uniqueness of one factor matrix of a CPD.

2.2. Uniqueness Conditions for CPD. A CPD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ is said to be unique if all the triplets $\left(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}, \widehat{\mathbf{C}}\right)$ satisfying (2.1) are related via

$$\widehat{\mathbf{A}} = \mathbf{A}\mathbf{P}\Delta_{\mathbf{A}}, \quad \widehat{\mathbf{B}} = \mathbf{B}\mathbf{P}\Delta_{\mathbf{B}}, \quad \widehat{\mathbf{C}} = \mathbf{C}\mathbf{P}\Delta_{\mathbf{C}},$$

where $\Delta_{\mathbf{A}}$, $\Delta_{\mathbf{B}}$ and $\Delta_{\mathbf{C}}$ are diagonal matrices satisfying $\Delta_{\mathbf{A}}\Delta_{\mathbf{B}}\Delta_{\mathbf{C}} = \mathbf{I}_R$ and $\mathbf{P}$ is a permutation matrix.

Together with related results in [9], Theorem 2.2 below is one of the most relaxed and yet quite easy-to-check deterministic CPD uniqueness conditions that have been reported in the literature.

THEOREM 2.2. *Consider the PD of* $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ *in* (2.1). *If*

$$\begin{cases} \min\left(k_{\boldsymbol{A}}, k_{\boldsymbol{B}}\right) + k_{\boldsymbol{C}} \geq R+1, \\ \max\left(k_{\boldsymbol{A}}, k_{\boldsymbol{B}}\right) + k_{\boldsymbol{C}} \geq R+2, \\ C_{R-r_{\boldsymbol{C}}+2}\left(\boldsymbol{A}\right) \odot C_{R-r_{\boldsymbol{C}}+2}\left(\boldsymbol{B}\right) \ \textit{ has full column rank,} \end{cases} \tag{2.6}$$

then the rank of $\mathcal{X}$ *is* R *and the CPD of* $\mathcal{X}$ *is unique [9].*

3. CPD with known factor matrix. Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ with matrix representation $\mathbf{X}_{(1)} = (\mathbf{A} \odot \mathbf{B})\,\mathbf{C}^T$. Assume that the factor matrix $\mathbf{C}$ is known and let the pair $\left(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}\right)$ yield an alternative decomposition of $\mathcal{X}$ with the same $\mathbf{C}$. The CPD of $\mathcal{X}$ with the known factor matrix $\mathbf{C}$ is said to be unique if all the pairs $\left(\widehat{\mathbf{A}}, \widehat{\mathbf{B}}\right)$ satisfying (2.1) are related via

$$\widehat{\mathbf{A}} = \mathbf{A}\Delta_{\mathbf{A}}\,, \quad \widehat{\mathbf{B}} = \mathbf{B}\Delta_{\mathbf{B}}\,,$$

where $\Delta_{\mathbf{A}}$ and $\Delta_{\mathbf{B}}$ are diagonal matrices with property $\Delta_{\mathbf{A}}\Delta_{\mathbf{B}} = \mathbf{I}_R$.

If the known factor matrix has full column rank, then the CPD with a known factor is unique in a trivial manner and can be computed via a number of rank-1 approximations. We state this result here for completeness.

PROPOSITION 3.1. *Consider the PD of* $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ *in* (2.1). *If* $\boldsymbol{C}$ *is known and has full column rank then the CPD of* $\mathcal{X}$ *with* $\boldsymbol{C}$ *known is unique (e.g. [28, 6]).*

Proof. Simply compute $\mathbf{Y} = \mathbf{X}_{(1)}\left(\mathbf{C}^T\right)^{\dagger}$, then the factor matrices are obtained by recognizing that the columns of $\mathbf{Y}$ are vectorized rank-1 matrices:

$$\min_{\mathbf{a}_r, \mathbf{b}_r} \left\|\mathbf{y}_r - \mathbf{a}_r \otimes \mathbf{b}_r\right\|_F^2\,, \quad r \in \{1, \ldots, R\}\,.$$

□

The following Proposition 3.2 presents a uniqueness condition for the case where the known factor matrix is not required to have full column rank.

PROPOSITION 3.2. *Consider the PD of* $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ *in* (2.1). *If* $\boldsymbol{C}$ *is known and*

$$k_{\boldsymbol{C}} + \min\left(\min\left(k_{\boldsymbol{A}}, k_{\boldsymbol{B}}\right) - 1, \max\left(k_{\boldsymbol{A}}, k_{\boldsymbol{B}}\right) - 2\right) \geq R, \tag{3.1}$$

then the CPD of $\mathcal{X}$ *with* $\boldsymbol{C}$ *known is unique [9].*

Except for the case where $\mathbf{C}$ has full column rank and $\max\left(k_{\mathbf{A}}, k_{\mathbf{B}}\right) = 1$, Proposition 3.2 yields a more relaxed condition than Proposition 3.1.

4. New uniqueness result for CPD with a known factor matrix. In subsection 4.1 we present a new result for the case where the known factor matrix does not necessarily have full column rank, but one of the unknown factors has. Subsection 4.2 generalizes the result to the case where none of the involved factor matrices is required to have full column rank.

4.1. At least one factor matrix has full column rank. The main result of this subsection is stated in Theorem 4.6. We first provide some auxiliary results.

LEMMA 4.1. *Given an analytic function* $f : \mathbb{C}^n \to \mathbb{C}$. *If there exists a vector* $\boldsymbol{x} \in \mathbb{C}^n$ *such that* $f\left(\boldsymbol{x}\right) \neq 0$, *then the set* $\{\, \boldsymbol{x} \,|\, f\left(\boldsymbol{x}\right) = 0 \,\}$ *is of Lebesgue measure zero (e.g. [11]).*

A matrix $\mathbf{A} \in \mathbb{C}^{I \times R}$ is said to be Vandermonde if it takes the form

$$\mathbf{A} = [\mathbf{a}_1, \ldots, \mathbf{a}_R] \,, \quad \mathbf{a}_r = \left[1, z_r, z_r^2, \ldots, z_r^{I-1}\right]^T , \tag{4.1}$$

where $\{z_r\}$ are called the generators of $\mathbf{A}$. Furthermore, a matrix/tensor decomposition property is called generic if the involved parameters are randomly drawn from absolutely continuous probability density functions. Intuitively, a generic property holds almost-surely for sufficiently random data.

LEMMA 4.2. *Let $\boldsymbol{A} \in \mathbb{C}^{I \times R}$ be a Vandermonde matrix and let $\boldsymbol{B} \in \mathbb{C}^{J \times R}$, then the matrix $\boldsymbol{A} \odot \boldsymbol{B}$ generically has rank* $\min(IJ, R)$.

Proof. The result follows from a combination of Theorem 3 and Corollary 1 in [11]. □

LEMMA 4.3. *Given a Vandermonde matrix $\boldsymbol{A} = [\boldsymbol{a}_1, \ldots, \boldsymbol{a}_R] \in \mathbb{C}^{I \times R}$ with distinct generators. Let the column vectors of $\boldsymbol{U} \in \mathbb{C}^{IJ \times R}$ constitute a basis for range$(\boldsymbol{A} \odot \boldsymbol{B})$, where $\boldsymbol{B} \in \mathbb{C}^{J \times R}$. If the matrix $\underline{\boldsymbol{A}} \odot \boldsymbol{B}$ has full column rank, then the matrix*

$$\boldsymbol{G}^{(r)} = [\boldsymbol{U}, \boldsymbol{a}_r \otimes \boldsymbol{I}_J] \in \mathbb{C}^{IJ \times (R+J)} \tag{4.2}$$

has rank $R + J - 1$, $\forall r \in \{1, \ldots, R\}$. Generically, $\boldsymbol{G}^{(r)}$ has rank $R + J - 1$ if $(I-1)J + 1 \geq R$.

Proof. A proof is provided in the supplementary material. □

A consequence of Lemma 4.3 is the following result, which generalizes the generic rank condition to the case where $\mathbf{A}$ is not Vandermonde.

COROLLARY 4.4. *Given $\boldsymbol{A} = [\boldsymbol{a}_1, \ldots, \boldsymbol{a}_R] \in \mathbb{C}^{I \times R}$ and let the column vectors of $\boldsymbol{U} \in \mathbb{C}^{IJ \times R}$ constitute a basis for range$(\boldsymbol{A} \odot \boldsymbol{B})$ in which $\boldsymbol{B} \in \mathbb{C}^{J \times R}$. If $(I-1)J+1 \geq R$, then the matrix*

$$\boldsymbol{G}^{(r)} = [\boldsymbol{U}, \boldsymbol{a}_r \otimes \boldsymbol{I}_J] \in \mathbb{C}^{IJ \times (R+J)} \tag{4.3}$$

generically has rank $R + J - 1$, $\forall r \in \{1, \ldots, R\}$.

Proof. A proof is provided in the supplementary material. □

In the proof of Theorem 4.6 we also make use of the following result.

PROPOSITION 4.5. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I \times J \times K}$ in (2.1). Assume that the rth column vector $\boldsymbol{c}_r$ of the factor matrix $\boldsymbol{C}$ is known. If* [1]

$$\begin{cases} \boldsymbol{B} \text{ has full column rank,} \\ r\left([\boldsymbol{C} \odot \boldsymbol{A}, \boldsymbol{c}_r \otimes \boldsymbol{I}_I]\right) = I + R - 1 \,, \end{cases} \tag{4.4}$$

then the column vector $\boldsymbol{a}_r$ of is $\boldsymbol{A}$ unique. Generically, condition (4.4) is satisfied if

$$\min\left((K-1)\, I + 1, J\right) \geq R \,. \tag{4.5}$$

Proof. Let us first prove that the deterministic condition (4.4) guarantees the uniqueness of $\mathbf{a}_r$. Note that, trivially, $\mathbf{c}_r \otimes \mathbf{a}_r \in \text{range}\,(\mathbf{c}_r \otimes \mathbf{I}_I)$. Condition (4.4) implies that this is the only dependency between the column vectors of $[\mathbf{C} \odot \mathbf{A}, \mathbf{c}_r \otimes \mathbf{I}_I]$. Hence, we know that $r\,(\mathbf{C} \odot \mathbf{A}) = R$. Given are $\mathbf{X}_{(3)} = (\mathbf{C} \odot \mathbf{A})\,\mathbf{B}^T$ and $\mathbf{c}_r$. We know that, under condition (4.4), both the matrices $\mathbf{C} \odot \mathbf{A}$ and $\mathbf{B}$ have full column

[1] The last condition states that $\mathbf{M}_r = [\mathbf{C} \odot \mathbf{A}, \mathbf{c}_r \otimes \mathbf{I}_I]$ has a one-dimensional kernel, which is minimal since $[\mathbf{n}_r^T, \mathbf{a}_r^T]^T \in \ker(\mathbf{M}_r)$ for some $\mathbf{n}_r \in \mathbb{C}^R$.

rank. Let $\mathbf{X}_{(3)} = \mathbf{U}\Sigma\mathbf{V}^H$ denote the compact SVD of $\mathbf{X}_{(3)}$, where $\mathbf{U} \in \mathbb{C}^{KI\times R}$, then there exists a nonsingular matrix $\mathbf{M} \in \mathbb{C}^{R\times R}$ such that

$$\mathbf{UM} = \mathbf{C} \odot \mathbf{A} \Leftrightarrow \mathbf{U}\mathbf{m}_r = \mathbf{c}_r \otimes \mathbf{a}_r, \quad r \in \{1, \ldots, R\}\,. \tag{4.6}$$

Relation (4.6) can also be written as

$$\mathbf{G}^{(r)} \begin{bmatrix} \mathbf{m}_r \\ \mathbf{a}_r \end{bmatrix} = \mathbf{0}_{KI}\,, \quad r \in \{1, \ldots, R\}\,, \tag{4.7}$$

where

$$\mathbf{G}^{(r)} = [\mathbf{U}, -\mathbf{c}_r \otimes \mathbf{I}_I] \in \mathbb{C}^{KI\times(I+R)}. \tag{4.8}$$

Since we assume that $\mathbf{G}^{(r)}$ has a one-dimensional kernel, the rth column vector of $\mathbf{A}$ follows from (4.7). Indeed, let $\mathbf{y}_r \in \mathbb{C}^{I+R}$ denote the right singular vector associated with the zero singular value of the matrix in (4.8), then $\mathbf{a}_r = \mathbf{y}_r\,(R+1:R+I)$.

Let us now prove that condition (4.5) generically guarantees the uniqueness of the column vector $\mathbf{a}_r$. Given are $\mathbf{X}_{(3)} = (\mathbf{C} \odot \mathbf{A})\,\mathbf{B}^T$ and $\mathbf{c}_r$. If $J \geq R$, then $\mathbf{B}$ generically has full column rank. Due to Lemma 4.2 we also know that since $KI \geq R$, $\mathbf{C} \odot \mathbf{A}$ generically has full column rank. Finally, Corollary 4.4 tells us that if $(K-1)\,J+1 \geq R$, then the rank of $\mathbf{G}^{(r)}$ in (4.8) is generically equal to $I+R-1$. Hence, generically we can work as in the proof of the deterministic case. □

THEOREM 4.6. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in* (2.1)*. Assume that the factor matrix $\boldsymbol{C}$ is known. If*

$$\begin{cases} \boldsymbol{B} \text{ has full column rank,} \\ r\left([\boldsymbol{C} \odot \boldsymbol{A}, \boldsymbol{c}_r \otimes \boldsymbol{I}_I]\right) = I + R - 1\,,\ \forall r \in \{1, \ldots, R\}\,, \end{cases} \tag{4.9}$$

then the CPD of $\mathcal{X}$ with $\boldsymbol{C}$ known is unique. Generically, condition (4.9) *is satisfied if*

$$\min\left((K-1)\,I+1, J\right) \geq R\,. \tag{4.10}$$

Proof. Let us first prove that condition (4.9) guarantees the uniqueness of the CPD of $\mathcal{X}$ with $\mathbf{C}$ known. Proposition 4.5 together with the condition (4.9) guarantees the uniqueness of the factor matrix $\mathbf{A}$. Recall also from the proof of Proposition 4.5 that the matrix $\mathbf{C} \odot \mathbf{A}$ has full column rank when condition (4.9) holds. Hence, $\mathbf{B}$ follows from

$$\mathbf{B}^T = (\mathbf{C} \odot \mathbf{A})^{\dagger}\,\mathbf{X}_{(3)} = \left(\left(\mathbf{C}^H\mathbf{C}\right) * \left(\mathbf{A}^H\mathbf{A}\right)\right)^{-1}(\mathbf{C} \odot \mathbf{A})^H\,\mathbf{X}_{(3)}\,.$$

The proof that condition (4.10) generically guarantees the uniqueness of the CPD of $\mathcal{X}$ with $\mathbf{C}$ is analogous to the proof of the generic part of Proposition 4.5. □

An important remark concerning the proof of Theorem 4.6 is that it is constructive, i.e., it provides us with an algorithm to compute a CPD with a known factor. In other words, if one of the factor matrices of the CPD is known, say $\mathbf{C}$, and the conditions stated in Theorem 4.6 are satisfied, then even if the known factor matrix does not have full column rank (as opposed to Proposition 3.1), we can find the remaining factor matrices from $\mathbf{X}_{(3)}$. An outline of the proposed method for computing a CPD with a known factor matrix is given in Algorithm 1.

Algorithm 1 Computation of CPD with known factor matrix based on Theorem 4.6.

Input: $\mathbf{X}_{(3)} = (\mathbf{C} \odot \mathbf{A})\,\mathbf{B}^T$ and $\mathbf{C}$.
Find $\mathbf{U}$ whose column vectors constitute an orthonormal basis for range $(\mathbf{C} \odot \mathbf{A})$.
Solve set of linear equations $[\mathbf{U}, -\mathbf{c}_r \otimes \mathbf{I}_I]\,\mathbf{y}_r = \mathbf{0}_{KI}, \quad r \in \{1, \ldots, R\}$.
Set $\mathbf{a}_r = \mathbf{y}_r\,(R+1 : R+I), \quad r \in \{1, \ldots, R\}$.
Compute $\mathbf{B}^T = \left(\left(\mathbf{C}^H\mathbf{C}\right) * \left(\mathbf{A}^H\mathbf{A}\right)\right)^{-1} (\mathbf{C} \odot \mathbf{A})^H\,\mathbf{X}_{(3)}$.
Output: $\mathbf{A}$ and $\mathbf{B}$.

Note that Theorem 4.6 does not prevent $k_{\mathbf{A}} = 1$. This is a remarkable difference with unconstrained CPD, where $k_{\mathbf{A}} \geq 2$ is necessary for uniqueness (e.g. [26]). Note also that cases where $\mathbf{C}$ does not have full column rank and $k_{\mathbf{A}} = 1$ are not covered by Propositions 3.1 and 3.2. More precisely, if $r_{\mathbf{C}} < R$, Proposition 3.1 does not apply. Likewise, if $k_{\mathbf{C}} < R$ and $k_{\mathbf{A}} = 1$, Proposition 3.1 does not apply either. Theorem 4.6 also leads to improved generic bounds. As an example, let $I = 4$, $J = R$, $K = 3$. Proposition 3.1 generically requires that $R \leq K = 3$ while Proposition 3.1 generically requires that $R \leq 6$. Theorem 4.6 relaxes the generic bound to $R \leq 9$.

4.2. None of the factor matrices is required to have full column rank. In Theorem 4.8 we explain that by an appropriate preprocessing step it is possible to derive a relaxed version of Theorem 4.6 in which $\mathbf{A}$ nor $\mathbf{B}$ are required to have full column rank. Proposition 4.7 below generalizes Proposition 4.5 to the case where none of the factor matrices is required to have full column rank. Proposition 4.7 makes use of the subsets S, S^c, T and U of $\{1, \ldots, R\}$. They satisfy the relations $S \subseteq T \subseteq \{1, \ldots, R\}$, $S^c = \{1, \ldots, R\} \setminus S$ and $U = T \setminus S \subseteq S^c$. The relations among the sets are visualized in the following figure 4.1.



FIG. 4.1. *The upper line depicts the disjoint partitioning of the set $\{1, \ldots, R\}$ into S and S^c. The middle line depicts the known columns of $\boldsymbol{C}$ indexed by the elements in the set T. The bottom line depicts the columns of $\boldsymbol{A}$ indexed by the elements in the set $U = T \setminus S$ for which uniqueness will be established in Proposition 4.7.*

PROPOSITION 4.7. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I \times J \times K}$ in (2.1). Assume that the columns of $\boldsymbol{C}$ indexed by the elements of the set T with $card(T) = Q$ are known. Stack the columns of $\boldsymbol{C}$ with index in T in $\boldsymbol{C}^{(T)} \in \mathbb{C}^{K \times card(T)}$. Let S denote a subset of T and let $S^c = \{1, \ldots, R\} \setminus S$. Stack the columns of $\boldsymbol{C}$ with index in S in $\boldsymbol{C}^{(S)} \in \mathbb{C}^{K \times card(S)}$ and stack the columns of $\boldsymbol{C}$ with index in S^c in $\boldsymbol{C}^{(S^c)} \in \mathbb{C}^{K \times (R - card(S))}$. Stack the columns of $\boldsymbol{A}$ (resp. $\boldsymbol{B}$) in the same order such that $\boldsymbol{A}^{(S)} \in \mathbb{C}^{I \times card(S)}$ (resp. $\boldsymbol{B}^{(S)} \in \mathbb{C}^{J \times card(S)}$) and $\boldsymbol{A}^{(S^c)} \in \mathbb{C}^{I \times (R - card(S))}$ (resp. $\boldsymbol{B}^{(S^c)} \in \mathbb{C}^{J \times (R - card(S))}$)*

are obtained. If there exists a subset $S \subseteq T$ *with* $0 \leq card\,(S) \leq r_{\boldsymbol{C}^{(T)}}$ *such that* [2]

$$\begin{cases} \boldsymbol{C}^{(S)} \text{ has full column rank,} \\ \boldsymbol{B}^{(S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{C}^{(S^c)}\right) \odot \boldsymbol{A}^{(S^c)}, (\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{c}_r) \otimes \boldsymbol{I}_I\right]\right) = I + R - card\,(S) - 1,\ \forall r \in U, \end{cases} \tag{4.11}$$

where $U = T \backslash S \subseteq S^c$*, then the column vectors* $\{\boldsymbol{a}_r\}_{r\in U}$ *of* $\boldsymbol{A}$ *are unique. Generically, condition* (4.11) *is satisfied if*

$$\begin{cases} R \leq \min\left(J + \min\left(K, Q\right), \frac{J(I-1)+I(K-1)+1}{I}\right) & when \quad J < R\,, \\ R \leq (K-1)I + 1 & when \quad J \geq R\,. \end{cases} \tag{4.12}$$

Proof. Let us first prove the deterministic part of the proposition. W.l.o.g. we assume that $\mathbf{C}(1 : \mathrm{card}\,(S), 1 : \mathrm{card}\,(S))$ is nonsingular, i.e., we set $\mathbf{C}^{(S)} = \mathbf{C}(:, 1 : \mathrm{card}\,(S))$. We first project on the orthogonal complement of range $\left(\mathbf{C}^{(S)}\right)$ to cancel $\mathrm{card}\,(S)$ rank-1 terms in the CPD of $\mathcal{X}$. That is, we compute

$$\mathbf{Y}_{(1)} = \mathbf{X}_{(1)}\mathbf{P}^T_{\mathbf{C}^{(S)}} = \left(\mathbf{A}^{(S^c)} \odot \mathbf{B}^{(S^c)}\right)\mathbf{C}^{(S^c)T}\mathbf{P}^T_{\mathbf{C}^{(S)}},$$

where the relation $\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C} = \mathbf{P}_{\mathbf{C}^{(S)}}\left[\mathbf{C}^{(S)}, \mathbf{C}^{(S^c)}\right] = \left[\mathbf{0}_{K,\mathrm{card}(S)}, \mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right]$ was used. The tensor $\mathcal{Y}$ represented by $\mathbf{Y}_{(1)}$ also has matrix representation

$$\mathbf{Y}_{(3)} = \left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)} \odot \mathbf{A}^{(S^c)}\right)\mathbf{B}^{(S^c)T}.$$

We have assumed that $r\left(\left[\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}, (\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r) \otimes \mathbf{I}_I\right]\right) = I + R - \mathrm{card}\,(S) - 1$. This implies that $r\left(\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}\right) = R - \mathrm{card}\,(S)$, i.e., $\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}$ has full column rank. We have also assumed that $\mathbf{B}^{(S^c)}$ has full column rank. Let $\mathbf{Y}_{(3)} = \mathbf{U}\Sigma\mathbf{V}^H$ denote the compact SVD of $\mathbf{Y}_{(3)}$ in which $\mathbf{U} \in \mathbb{C}^{KI\times(R-\mathrm{card}(S))}$, then there exists a nonsingular matrix $\mathbf{M} \in \mathbb{C}^{(R-\mathrm{card}(S))\times(R-\mathrm{card}(S))}$ such that

$$\mathbf{UM} = \left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}$$

and

$$\mathbf{G}^{(r,S^c)}\begin{bmatrix} \mathbf{m}_{\mu(r)} \\ \mathbf{a}_r \end{bmatrix} = \mathbf{0}_{KI}\,, \quad r \in S^c, \tag{4.13}$$

in which $\mathbf{M} = \left[\mathbf{m}_{\mu(1)}, \ldots, \mathbf{m}_{\mu(\mathrm{card}(S^c))}\right]$ and $\mathbf{G}^{(r,S^c)} \in \mathbb{C}^{KI\times(I+R-\mathrm{card}(S))}$ is given by

$$\mathbf{G}^{(r,S^c)} = \left[\mathbf{U}, -\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r\right) \otimes \mathbf{I}_I\right]\,, \quad r \in S^c. \tag{4.14}$$

We have assumed that $\mathbf{G}^{(r,S^c)}$ has rank $I + R - \mathrm{card}\,(S) - 1$, $\forall r \in U \subseteq S^c$, which implies that $\mathbf{G}^{(r,S^c)}$ has a one-dimensional kernel for every $r \in U$. This in turn

[2] The last condition states that $\mathbf{M}_r = \left[\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}, \left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r^{(S^c)}\right) \otimes \mathbf{I}_I\right]$ has a one-dimensional kernel for every $r \in U$, which is minimal since $[\mathbf{n}_r^T, \mathbf{a}_r^{(S^c)T}]^T \in \ker\left(\mathbf{M}_r\right)$ for some $\mathbf{n}_r \in \mathbb{C}^{\mathrm{card}(S^c)}$.

means that we can obtain the column vectors $\{\mathbf{a}_r\}_{r\in U}$ from (4.13) as follows. Let $\mathbf{z}_r \in \mathbb{C}^{(I+R-\mathrm{card}(S))}$ denote the right singular vector associated with the zero singular value of the matrix given by (4.14), then

$$\mathbf{a}_r = \mathbf{z}_r \left(R - \mathrm{card}\,(S) + 1 : R - \mathrm{card}\,(S) + I\right), \quad r \in U.$$

Let us now prove that condition (4.12) generically guarantees the uniqueness of the column vectors $\{\mathbf{a}_r\}_{r\in U}$.

Consider first the case where $J \geq R$. In that case we choose $S = \emptyset$ and consequently also $\mathbf{P}_{\mathbf{C}^{(S)}} = \mathbf{I}_K$. We know that $\mathbf{B}$ generically has full column rank when $J \geq R$. Due to Corollary 4.4 we also know that $[\mathbf{C} \odot \mathbf{A}, \mathbf{c}_r \otimes \mathbf{I}_I]$ generically has rank $R + I - 1$ when $(K-1)I + 1 \geq R$. This proves the second leg of (4.12).

Consider now the case where $J < R$. Due to Lemma 4.1 we only need to find one example for which the first leg of (4.12) generically guarantees the uniqueness of the column vectors $\{\mathbf{a}_r\}_{r\in U}$. By way of example, we let

$$\mathbf{C} = \left[\begin{array}{c|c} \mathbf{C}^{(S)} & \mathbf{C}^{(S^c)} \end{array}\right] = \left[\begin{array}{c|c} \mathbf{I}_{\mathrm{card}(S)} & \\ \mathbf{0}_{K-\mathrm{card}(S),\mathrm{card}(S)} & \mathbf{C}^{(S^c)} \end{array}\right]. \tag{4.15}$$

Then

$$\mathbf{P}_{\mathbf{C}^{(S)}} = \mathbf{I}_K - \left[\begin{array}{c|c} \mathbf{I}_{\mathrm{card}(S)} & \mathbf{0}_{\mathrm{card}(S),K-\mathrm{card}(S)} \\ \mathbf{0}_{K-\mathrm{card}(S),\mathrm{card}(S)} & \mathbf{0}_{K-\mathrm{card}(S),K-\mathrm{card}(S)} \end{array}\right]$$

and

$$\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)} = \left[\begin{array}{c} \mathbf{0}_{\mathrm{card}(S),R-\mathrm{card}(S)} \\ \mathbf{D} \end{array}\right], \quad \mathbf{D} \in \mathbb{C}^{(K-\mathrm{card}(S))\times(R-\mathrm{card}(S))}.$$

Note that $\mathrm{card}\,(S) = r_{\mathbf{C}^{(T)}} - m$ for some integer m with property $0 \leq m \leq r_{\mathbf{C}^{(T)}}$. We know that $\mathbf{B}^{(S^c)} \in \mathbb{C}^{J\times(R-\mathrm{card}(S))}$ generically has full column rank if

$$J \geq R - \mathrm{card}\,(S) = R - r_{\mathbf{C}^{(T)}} + m \Leftrightarrow J - R + r_{\mathbf{C}^{(T)}} \geq m. \tag{4.16}$$

Note that, since $J < R$, the condition $m \leq r_{\mathbf{C}^{(T)}}$ is automatically satisfied when (4.16) holds. In order to ensure that $m \geq 0$ the following inequality must be satisfied:

$$J + r_{\mathbf{C}^{(T)}} \geq R\,, \tag{4.17}$$

where $r_{\mathbf{C}^{(T)}} = \min(K, Q)$. Due to the structure of $\mathbf{C}$ in (4.15) the problem of determining the generic rank of

$$\left[\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}, (\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r) \otimes \mathbf{I}_I\right]$$

reduces to finding the generic rank of

$$\left[\mathbf{D} \odot \mathbf{A}^{(S^c)}, \mathbf{d}_r \otimes \mathbf{I}_I\right] \in \mathbb{C}^{(K-\mathrm{card}(S))I\times(R-\mathrm{card}(S)+I)}. \tag{4.18}$$

Corollary 4.4 tells us that the matrix given by (4.18) generically has rank $R-\mathrm{card}\,(S)+I-1$ if

$$(K - \mathrm{card}\,(S) - 1)I + 1 \geq R - \mathrm{card}\,(S)\,. \tag{4.19}$$

By inserting $\text{card}\,(S) = r_{\mathbf{C}^{(T)}} - m$ into (4.19) we obtain

$$(K - r_{\mathbf{C}^{(T)}} + m - 1)I + 1 \geq R - r_{\mathbf{C}^{(T)}} + m. \tag{4.20}$$

Relation (4.20) can also be expressed as

$$m \geq \frac{R + r_{\mathbf{C}^{(T)}}(I-1) - I(K-1) - 1}{I-1}. \tag{4.21}$$

Combining conditions (4.16) and (4.21) we conclude that if the inequalities (4.17) and

$$\frac{R + r_{\mathbf{C}^{(T)}}(I-1) - I(K-1) - 1}{I-1} \leq J - R + r_{\mathbf{C}^{(T)}} \Leftrightarrow R \leq \frac{J(I-1) + I(K-1) + 1}{I}$$

are satisfied, then generically the column vectors $\{\mathbf{a}_r\}_{r\in U}$ are unique. This proves the first leg of (4.12). □

Note that there is some flexibility in the choice of $\text{card}\,(S)$ in (4.11). By choosing a large $\text{card}\,(S)$ we relax the constraint on $\mathbf{B}^{(S^c)}$ but also impose a stronger constraint on $\mathbf{C}$. The larger $\text{card}\,(S)$, the smaller the number of columns of $\mathbf{A}$ for which uniqueness is demonstrated.

THEOREM 4.8. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in (2.1). Assume that $\boldsymbol{C}$ is known. Let S denote a subset of $\{1,\ldots,R\}$ and let $S^c = \{1,\ldots,R\} \setminus S$ denote the complementary set. Stack the columns of $\boldsymbol{C}$ with index in S in $\boldsymbol{C}^{(S)} \in \mathbb{C}^{K\times \text{card}(S)}$ and stack the columns of $\boldsymbol{C}$ with index in S^c in $\boldsymbol{C}^{(S^c)} \in \mathbb{C}^{K\times(R-\text{card}(S))}$. Stack the columns of $\boldsymbol{A}$ (resp. $\boldsymbol{B}$) in the same order such that $\boldsymbol{A}^{(S)} \in \mathbb{C}^{I\times \text{card}(S)}$ (resp. $\boldsymbol{B}^{(S)} \in \mathbb{C}^{J\times \text{card}(S)}$) and $\boldsymbol{A}^{(S^c)} \in \mathbb{C}^{I\times(R-\text{card}(S))}$ (resp. $\boldsymbol{B}^{(S^c)} \in \mathbb{C}^{J\times(R-\text{card}(S))}$) are obtained. If there exists a subset S of $\{1,\ldots,R\}$ with $0 \leq \text{card}(S) \leq r_{\boldsymbol{C}}$ such that $\boldsymbol{C}^{(S)}$ has full column rank and*

$$\begin{cases} \boldsymbol{B}^{(S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{C}^{(S^c)}\right) \odot \boldsymbol{A}^{(S^c)}, (\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{c}_r) \otimes \boldsymbol{I}_I\right]\right) = \alpha,\ \forall r \in S^c, \end{cases} \tag{4.22a}$$

where $\alpha = I + R - \text{card}(S) - 1$, or

$$\begin{cases} \boldsymbol{A}^{(S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{C}^{(S^c)}\right) \odot \boldsymbol{B}^{(S^c)}, (\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{c}_r) \otimes \boldsymbol{I}_J\right]\right) = \beta,\ \forall r \in S^c, \end{cases} \tag{4.22b}$$

where $\beta = J + R - \text{card}(S) - 1$, then the CPD of $\mathcal{X}$ with $\boldsymbol{C}$ known is unique. Generically, condition (4.22a) or (4.22b) is satisfied if

$$\begin{cases} R \leq \min\left(W + \min(K,R), \frac{W(V-1)+V(K-1)+1}{V}\right) & \text{when} \quad W < R, \\ R \leq (K-1)V + 1 & \text{when} \quad W \geq R, \end{cases} \tag{4.23}$$

where $V = \min(I,J)$ and $W = \max(I,J)$.

Proof. Let us first prove that condition (4.22a) guarantees the uniqueness of the CPD of $\mathcal{X}$ with $\mathbf{C}$ known. We work as follows. The fact that $\mathbf{C}$ is known allows us to project away the part of the CPD of $\mathcal{X}$ that corresponds to S. After finding $\mathbf{A}^{(S^c)}$ and $\mathbf{B}^{(S^c)}$, we subtract the part of the CPD of $\mathcal{X}$ that corresponds to S^c, which leads to the remaining $\mathbf{A}^{(S)}$ and $\mathbf{B}^{(S)}$.

More in detail, we first determine the matrix $\mathbf{A}^{(S^c)}$ in a similar way as the vectors $\{\mathbf{a}\}_{r\in U}$ in Proposition 4.7. More precisely, we set $Q = R$ and consequently $T = \{1,\ldots,R\}$, $U = \{1,\ldots,R\} \setminus S = S^c$.

The next step is to find $\mathbf{B}^{(S^c)}$. Recall that we assumed that

$$r\left(\left[\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}, (\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r) \otimes \mathbf{I}_I\right]\right) = I + R - \operatorname{card}(S) - 1.$$

This implies that $r\left(\left(\mathbf{P}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}\right) = R - \operatorname{card}(S)$, i.e., $\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}$ has full column rank. Hence, $\mathbf{B}^{(S^c)}$ follows from the relation

$$\begin{aligned}\mathbf{B}^{(S^c)T} &= \left(\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}\right)^{\dagger} \mathbf{Y}_{(3)} \\ &= \left(\left(\mathbf{C}^{(S^c)H}\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) * \left(\mathbf{A}^{(S^c)H}\mathbf{A}^{(S^c)}\right)\right)^{-1} \left(\mathbf{P}\mathbf{C}^{(S^c)} \odot \mathbf{A}^{(S^c)}\right)^{H} \mathbf{Y}_{(3)}\,.\end{aligned}$$

Now that $\left\{\mathbf{A}^{(S^c)}, \mathbf{B}^{(S^c)}, \mathbf{C}^{(S^c)}\right\}$ are known, we can compute

$$\mathbf{Q}_{(1)} = \mathbf{X}_{(1)} - \left(\mathbf{A}^{(S^c)} \odot \mathbf{B}^{(S^c)}\right)\mathbf{C}^{(S^c)T} = \left(\mathbf{A}^{(S)} \odot \mathbf{B}^{(S)}\right)\mathbf{C}^{(S)T}.$$

Recall also that the matrix $\mathbf{C}^{(S)}$ is assumed to have full column rank. Compute

$$\mathbf{H} = \mathbf{Q}_{(1)}\left(\mathbf{C}^{(S)T}\right)^{\dagger}.$$

The remaining unknowns columns of $\mathbf{A}^{(S)}$ and $\mathbf{B}^{(S)}$ are obtained by recognizing that the columns of $\mathbf{H}$ are vectorized rank-1 matrices:

$$\min_{\mathbf{a}_r,\mathbf{b}_r} \left\|\mathbf{h}_{\sigma(r)} - \mathbf{a}_r \otimes \mathbf{b}_r\right\|_F^2\,, \quad r \in S\,,$$

where $\mathbf{H} = \left[\mathbf{h}_{\sigma(1)},\ldots,\mathbf{h}_{\sigma(S)}\right]$. The proof for (4.22b) is analogous to the above proof for condition (4.22a).

The proof that condition (4.23) generically guarantees the uniqueness of the CPD of $\mathcal{X}$ with $\mathbf{C}$ known is analogous to the generic part of the proof of Proposition 4.7. Essentially, by replacing Q with R in the proof of the generic condition (4.12) in Proposition 4.7 the result follows. □

Note that Theorem 4.8 does not prevent that $k_{\mathbf{A}} = 1$ and/or $k_{\mathbf{B}} = 1$ and/or $r_{\mathbf{A}\odot\mathbf{B}} < R$. This is in contrast with unconstrained CPD where $k_{\mathbf{A}} \geq 2$, $k_{\mathbf{B}} \geq 2$ and $r_{\mathbf{A}\odot\mathbf{B}} = R$ are all necessary uniqueness conditions (e.g. [26]). As an example, consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in (2.1) in which $\mathbf{C}$ is known, $\mathbf{a}_2 = \mathbf{a}_1$, $\mathbf{b}_2 = \mathbf{b}_1$, $R = 7$, $I = 4$, $J = 4$ and $K = 5$. If apart from these constraints the parameters are randomly drawn, Theorem 4.8 with $S = \{1,2\}$ and $\operatorname{card}(S) = 2$ guarantees the generic uniqueness of $\mathbf{A}$ and $\mathbf{B}$.

Theorem 4.8 also leads to improved bounds on R in generic cases without collinearities. As an example, we consider the situation where $V = \min(I,J) = 3$, $W = \max(I,J) = 7$, $K = 4$ and $\mathbf{C}$ is known. Proposition 3.1 requires that $R \leq K = 4$ while Proposition 3.2 relaxes the bound to $R \leq 6$. On the other hand, Theorem 4.8 only requires that $R \leq 8$.

We observe that the proof of Theorem 4.8 is constructive. We present the construction as Algorithm 2.

The cardinality of set S in Algorithm 2 must be chosen such that the conditions in Theorem 4.8 are satisfied. As explained in the proof, the choice of $\text{card}\,(S)$ affects the computation in the sense that we first compute the rank-$\text{card}\,(S^c)$ CPD $\mathcal{Y} = \sum_{r\in S^c} \mathbf{a}_r \circ \mathbf{b}_r \circ \mathbf{c}_r$ and thereafter the rank-$\text{card}\,(S)$ CPD $\mathcal{Q} = \sum_{r\in S} \mathbf{a}_r \circ \mathbf{b}_r \circ \mathbf{c}_r$. S must be chosen such that $\mathbf{C}^{(S)}$ has full column rank, which is easy to check. S must also be chosen such that $\mathbf{B}^{(S^c)}$ has full column rank, implying that $\text{card}\,(S) \geq R - J$ must be satisfied. This condition can numerically be verified by checking if the effective rank of $\mathbf{Y}_{(3)}$ is equal to $\text{card}\,(S^c)$. S must also be chosen such that $\left[\left(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}\right) \odot \mathbf{A}^{(S^c)}, (\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r) \otimes \mathbf{I}_I\right]$ has a one-dimensional kernel for every $r \in S^c$. This condition can numerically be verified by checking if the effective dimension of the kernel of $[\mathbf{U}, -(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r) \otimes \mathbf{I}_I]$ is one for every $r \in S^c$.

In the supplementary material we briefly explain how to extend Theorem 4.8 and Algorithm 2 to tensors of arbitrary order.

Algorithm 2 Computation of CPD with known factor matrix based on Theorem 4.8.

Input: $\mathbf{X}_{(1)} = (\mathbf{A} \odot \mathbf{B})\,\mathbf{C}^T$ and $\mathbf{C}$

1. Determine $r_{\mathbf{C}}$.
2. Choose sets $S \subseteq \{1, \ldots, R\}$ and $S^c = \{1, \ldots, R\} \backslash S$ subject to $0 \leq \text{card}\,(S) \leq r_{\mathbf{C}}$.
3. Build $\mathbf{C}^{(S)}$ and $\mathbf{C}^{(S^c)}$.
4. Find $\mathbf{F}$ whose column vectors constitute an orthonormal basis for range $\left(\mathbf{C}^{(S)}\right)$.
5. Compute $\mathbf{P}_{\mathbf{C}^{(S)}} = \mathbf{I}_K - \mathbf{F}\mathbf{F}^H$, $\mathbf{D}^{(S^c)} = \mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}$ and $\mathbf{E}^{(S^c)} = \mathbf{C}^{(S^c)H}\mathbf{D}^{(S^c)}$.
6. Compute $\mathbf{Y}_{(1)} = \mathbf{X}_{(1)}\mathbf{P}^T_{\mathbf{C}^{(S)}}$.
7. Build $\mathbf{Y}_{(3)}$.
8. Find $\mathbf{U}$ whose column vectors constitute an orthonormal basis for range $\left(\mathbf{Y}_{(3)}\right)$.
9. Solve set of linear equations $[\mathbf{U}, -(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r) \otimes \mathbf{I}_I]\,\mathbf{z}_r = \mathbf{0}_{KI}, \quad r \in S^c$.
10. Set $\mathbf{a}_r = \mathbf{z}_r\,(R - \text{card}\,(S) + 1 : R - \text{card}\,(S) + I)\,, \quad r \in S^c\,.$
11. Compute $\mathbf{B}^{(S^c)T} = \left(\mathbf{E}^{(S^c)} * \left(\mathbf{A}^{(S^c)H}\mathbf{A}^{(S^c)}\right)\right)^{-1}\left(\mathbf{D}^{(S^c)} \odot \mathbf{A}^{(S^c)}\right)^H \mathbf{Y}_{(3)}$.
12. Compute $\mathbf{Q}_{(1)} = \mathbf{X}_{(1)} - \left(\mathbf{A}^{(S^c)} \odot \mathbf{B}^{(S^c)}\right)\mathbf{C}^{(S^c)T}$.
13. Compute $\mathbf{H} = \left[\mathbf{h}_{\sigma(1)}, \ldots, \mathbf{h}_{\sigma(S)}\right] = \mathbf{Q}_{(1)}\left(\mathbf{C}^{(S)T}\right)^{\dagger}$.
14. Compute best rank-1 approximations $\min\limits_{\mathbf{a}_r, \mathbf{b}_r} \left\|\mathbf{h}_{\sigma(r)} - \mathbf{a}_r \otimes \mathbf{b}_r\right\|_F^2\,, \quad r \in S\,.$

Output: $\mathbf{A}$ and $\mathbf{B}$.

5. New uniqueness results for overall CPD. By combining the existing result presented in subsection 2.1 with the new results presented in section 4 we will in this section derive new uniqueness conditions for the overall CPD and also some variants that are important in signal processing.

5.1. New uniqueness condition for CPD. We first present a deterministic uniqueness condition for CPD based on Theorem 4.8.

THEOREM 5.1. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in (2.1). Let S denote a subset of $\{1, \ldots, R\}$ and let $S^c = \{1, \ldots, R\} \setminus S$ denote the complementary set. Stack the columns of $\boldsymbol{C}$ with index in S in $\boldsymbol{C}^{(S)} \in \mathbb{C}^{K\times card(S)}$ and stack the columns of $\boldsymbol{C}$ with index in S^c in $\boldsymbol{C}^{(S^c)} \in \mathbb{C}^{K\times(R-card(S))}$. Stack the columns of $\boldsymbol{A}$ (resp. $\boldsymbol{B}$) in the same order such that $\boldsymbol{A}^{(S)} \in \mathbb{C}^{I\times card(S)}$ (resp. $\boldsymbol{B}^{(S)} \in \mathbb{C}^{J\times card(S)}$) and $\boldsymbol{A}^{(S^c)} \in$*

$\mathbb{C}^{I\times(R-card(S))}$ *(resp.* $\boldsymbol{B}^{(S^c)} \in \mathbb{C}^{J\times(R-card(S))}$*) are obtained. If*

$$C_{R-r_{\boldsymbol{C}}+2}(\boldsymbol{A}) \odot C_{R-r_{\boldsymbol{C}}+2}(\boldsymbol{B}) \text{ has full column rank,} \tag{5.1a}$$

and if there exists a subset $S \subseteq \{1,\ldots,R\}$ *with* $0 \leq card(S) \leq r_{\boldsymbol{C}}$ *such that* $\boldsymbol{C}^{(S)}$ *has full column rank and*

$$\begin{cases} \boldsymbol{B}^{(S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{C}^{(S^c)}\right) \odot \boldsymbol{A}^{(S^c)}, (\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{c}_r^{(S^c)}) \otimes \boldsymbol{I}_I\right]\right) = \alpha, \ \forall r \in S^c, \end{cases} \tag{5.1b}$$

where $\alpha = I + R - card(S) - 1$*, or*

$$\begin{cases} \boldsymbol{A}^{(S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{C}^{(S^c)}\right) \odot \boldsymbol{B}^{(S^c)}, (\boldsymbol{P}_{\boldsymbol{C}^{(S)}}\boldsymbol{c}_r^{(S^c)}) \otimes \boldsymbol{I}_J\right]\right) = \beta, \ \forall r \in S^c, \end{cases} \tag{5.1c}$$

where $\beta = J + R - card(S) - 1$*, then the rank of* $\mathcal{X}$ *is* R *and the CPD of* $\mathcal{X}$ *is unique.*

Proof. Note that both conditions (5.1b) and (5.1c) imply that $k_{\mathbf{C}} \geq 2$. Indeed, if $k_{\mathbf{C}} = 1$, then (i) $k_{\mathbf{C}^{(S)}} = 1$, (ii) $k_{\mathbf{C}^{(S^c)}} = 1$ or (iii) $k_{[\mathbf{c}_m,\mathbf{c}_n]} = 1$ for some $m \in S$ and $n \in S^c$. Since it is assumed that $\mathbf{C}^{(S)}$ has full column rank (i.e., $k_{\mathbf{C}^{(S)}} = \text{card}(S)$) we can rule out the first case (i). The second and third cases imply that the dimension of the kernel of $[(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}) \odot \mathbf{A}^{(S^c)}, (\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r^{(S^c)}) \otimes \mathbf{I}_I]$ or $[(\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}) \odot \mathbf{B}^{(S^c)}, (\mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{c}_r^{(S^c)}) \otimes \mathbf{I}_J]$ is larger than one for some $r \in S^c$. Hence, cases (ii) and (iii) can also be ruled out. Theorem 2.1 together with the inequality $k_{\mathbf{C}} \geq 2 > 1$ now tells us that the rank of $\mathcal{X}$ is R and that the factor $\mathbf{C}$ is unique. Let us now consider $\mathcal{X}$ as a third-order tensor with known $\mathbf{C}$. Due to Theorem 4.8, the conditions (5.1b)/(5.1c) now guarantee the overall uniqueness of the CPD. □

For a discussion of Theorem 5.1, let us first consider the generic case, where factor matrices have maximal rank and k-rank. We notice that, generically, if the one factor matrix uniqueness condition in Theorem 2.1 is satisfied, then the overall uniqueness condition in Theorem 2.2 is automatically satisfied. This means that in the generic case, the one factor matrix uniqueness condition in Theorem 2.1, which corresponds to condition (5.1a), makes that Theorem 5.1 is not more relaxed than Theorem 2.2.

In order to demonstrate that Theorem 5.1 can nevertheless improve Theorem 2.2 in non-generic cases we consider the following example taken from [9]. Consider the third-order tensor $\mathcal{X} \in \mathbb{C}^{4\times4\times4}$ with matrix representation $\mathbf{X}_{(1)} = (\mathbf{A} \odot \mathbf{B})\,\mathbf{C}^T$ in which $\mathbf{A} \in \mathbb{C}^{4\times5}$, $\mathbf{B} \in \mathbb{C}^{4\times5}$ and $\mathbf{C} \in \mathbb{C}^{4\times5}$, where $[\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4] = \mathbf{I}_4$ and $a_{25} = 0$, $[\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_4] = \mathbf{I}_4$ and $b_{15} = 0$, and $[\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3, \mathbf{c}_4] = \mathbf{I}_4$ and $c_{25} = 0$. For a random choice of parameter entries we have $k_{\mathbf{A}} = 3$, $k_{\mathbf{B}} = 3$ and $k_{\mathbf{C}} = 3$. On the other hand, we also have $r_{\mathbf{A}} = 4$, $r_{\mathbf{B}} = 4$ and $r_{\mathbf{C}} = 4$. For this problem the conditions in Theorem 2.2 are not satisfied. However, the conditions (5.1a)–(5.1b) in Theorem 5.1 are generically satisfied if $\text{card}(S) = 2$ and $S = \{1, 2\}$. Thus, overall uniqueness of the CPD of $\mathcal{X}$ can be established by Theorem 5.1 but not by the related Theorem 2.2. We mention that CPD uniqueness of $\mathcal{X}$ has already been established in [9] by other means. A nice property of Theorem 5.1 is that it can be extended to other tensor decompositions than the third-order CPD. For instance, in [24, 25] we demonstrate how Theorem 5.1 can be adapted to coupled CPD and CPD of tensors of arbitrary order. As a final remark, we recall that Theorem 2.1 is not necessary for one factor uniqueness. For this reason, it may be possible to further generalize Theorem 5.1 by combining Theorem 4.8 with a more relaxed single factor matrix uniqueness condition.

5.2. New uniqueness condition for CPD with partial Hermitian symmetry. We say that the CPD of a tensor $\mathcal{X} \in \mathbb{C}^{I\times I\times K}$ has partial Hermitian symmetry if $\mathbf{X}_{(1)} = (\mathbf{A} \odot \mathbf{A}^*)\,\mathbf{C}^T$. The Hermitian symmetry-constrained rank of $\mathcal{X}$ is equal to the minimal number of rank-1 terms $\mathbf{a}_r \circ \mathbf{a}_r^* \circ \mathbf{c}_r$ that yield $\mathcal{X}$ in a linear combination. This structure is very common in signal separation applications. For instance, tensors with partial Hermitian symmetry are obtained by stacking sets of covariance matrices of complex data, see [2, 7] for references to concrete applications. A uniqueness condition for the Hermitian symmetry-constrained CPD of a tensor was provided in [27] for the special case where $\mathbf{C}$ has full column rank. We will now provide a new uniqueness condition. In contrast to [27] we do not require that $\mathbf{C}$ has full column rank. In fact, by better exploiting the Hermitian symmetry, $\mathbf{C}$ may even contain collinear columns and even $K = 1$ can be admitted in some cases. Let us define the tensor $\mathcal{Y} \in \mathbb{C}^{I\times I\times K}$ by $y_{ijk} = x_{jik}^*$ for all indices. We have $\mathcal{Z} \in \mathbb{C}^{I\times I\times 2K}$ with matrix representation

$$\mathbf{Z}_{(1)} = \left[\mathbf{X}_{(1)}, \mathbf{Y}_{(1)}\right] = (\mathbf{A} \odot \mathbf{A}^*)\,\mathbf{D}^T \in \mathbb{C}^{I^2\times 2K}, \tag{5.2}$$

where

$$\mathbf{D} = \begin{bmatrix} \mathbf{C} \\ \mathbf{C}^* \end{bmatrix} \in \mathbb{C}^{2K\times R}. \tag{5.3}$$

We may now study the uniqueness of the CPD of $\mathcal{Z}$ in order to obtain uniqueness results for $\mathcal{X}$. From relation (5.2) it is clear that if $\mathbf{A}$ has full column rank, then the Hermitian symmetry-constrained CPD of $\mathcal{X}$ is unique if $k_{\mathbf{D}} \geq 2$. Note that the latter condition does not prevent $k_{\mathbf{C}} = 1$ or even $K = 1$. Theorem 5.2 is an adaption of Theorem 5.1 to the Hermitian symmetry-constrained CPD case.

THEOREM 5.2. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times I\times K}$ with matrix representation $\boldsymbol{X}_{(1)} = (\boldsymbol{A} \odot \boldsymbol{A}^*)\,\boldsymbol{C}^T$. Consider also $\boldsymbol{D}$ in (5.3). Let S denote a subset of $\{1,\dots,R\}$ and let $S^c = \{1,\dots,R\} \setminus S$ denote the complementary set. Stack the columns of $\boldsymbol{D}$ with index in S in $\boldsymbol{D}^{(S)} \in \mathbb{C}^{2K\times card(S)}$ and stack the columns of $\boldsymbol{D}$ with index in S^c in $\boldsymbol{D}^{(S^c)} \in \mathbb{C}^{2K\times(R-card(S))}$. Stack the columns of $\boldsymbol{A}$ in the same order such that $\boldsymbol{A}^{(S)} \in \mathbb{C}^{I\times card(S)}$ and $\boldsymbol{A}^{(S^c)} \in \mathbb{C}^{I\times(R-card(S))}$ are obtained. If*

$$C_{R-r_{\boldsymbol{D}}+2}\left(\boldsymbol{A}^*\right) \odot C_{R-r_{\boldsymbol{D}}+2}\left(\boldsymbol{A}\right) \text{ has full column rank,} \tag{5.4a}$$

and if there exists a subset $S \subseteq \{1,\dots,R\}$ with $0 \leq card(S) \leq r_{\boldsymbol{D}}$ such that

$$\begin{cases} \boldsymbol{D}^{(S)} \text{ has full column rank,} \\ \boldsymbol{A}^{(S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{D}^{(S)}}\boldsymbol{D}^{(S^c)}\right) \odot \boldsymbol{A}^{(S^c)}, \left(\boldsymbol{P}_{\boldsymbol{D}^{(S)}}\boldsymbol{d}_r^{(S^c)}\right) \otimes \boldsymbol{I}_I\right]\right) = \alpha, \forall r \in S^c, \end{cases} \tag{5.4b}$$

where $\alpha = I + R - card(S) - 1$, then the Hermitian symmetry-constrained rank of $\mathcal{X}$ is R and the Hermitian symmetry-constrained CPD of $\mathcal{X}$ is unique.

Proof. Analogous to the proof of Theorem 5.1, condition (5.4b) implies that $k_{\mathbf{D}} \geq 2$. Consequently, conditions (5.4a) and (5.4b) together with Theorem 2.1 imply that the rank of $\mathcal{Z}$ with matrix representation (5.2) is R and that the factor $\mathbf{D}$ is unique. Since $\mathbf{D}$ is unique, $\mathbf{C}$ is unique. Due to Theorem 4.8, we know that if condition (5.4b) is satisfied, $\mathbf{A}$ is also unique. Overall, we obtain that the Hermitian symmetry-constrained CPD of $\mathcal{X}$ is unique and the Hermitian symmetry-constrained rank of $\mathcal{X}$ is R. ∎

Equations (5.2)–(5.3) are the key to understanding why partial Hermitian symmetry allows us to relax uniqueness conditions. Indeed, a condition on $\mathbf{D}$ is typically less restrictive than a condition on $\mathbf{C}$ (Note that generically $r_{\mathbf{C}} = \min(K, R)$, while, on the other hand, generically $r_{\mathbf{D}} = \min(2K, R)$, see the supplementary material). For that reason the conditions in Theorem 5.2 are more relaxed than the conditions for unconstrained CPD in Theorems 2.2 and 5.1. For instance, Theorem 5.2 does not prevent $k_{\mathbf{C}} = 1$, i.e., it only requires that $k_{\mathbf{D}} \geq 2$. This is in contrast with the unconstrained CPD for which $k_{\mathbf{C}} \geq 2$ is a necessary uniqueness condition (e.g. [26]). Theorem 5.2 is also more relaxed than the existing result presented in [27]. In contrast to [27], Theorem 5.2 allows us to establish uniqueness in cases where $r_{\mathbf{C}} < R$, $k_{\mathbf{C}} = 1$ and even $K = 1$.

A necessary condition for Hermitian symmetry-constrained CPD uniqueness is that the matrix $\mathbf{A} \odot \mathbf{A}^*$ must have full column rank. Indeed, if $\mathbf{A} \odot \mathbf{A}^*$ does not have full column rank, then for any $\mathbf{x} \in \ker(\mathbf{A} \odot \mathbf{A}^*)$ we obtain from relation (5.2) that $\mathbf{Z}_{(1)} = (\mathbf{A} \odot \mathbf{A}^*)\,\mathbf{D}^T = (\mathbf{A} \odot \mathbf{A}^*)\left(\mathbf{D}^T + \mathbf{x}\left[\mathbf{y}^T, \mathbf{y}^T\right]\right)$, where $\mathbf{y} \in \mathbb{R}^K$. Since $\mathbf{A} \odot \mathbf{A}^*$ generically has rank $\min(I^2, R)$ [23], this requirement is not very restrictive.

5.3. New uniqueness condition for CPD with columnwise orthonormal factor matrix. In this section we consider the PD of $\mathcal{X} \in \mathbb{C}^{I_1 \times I_2 \times K}$ with matrix representation $\mathbf{X}_{(1)} = \left(\mathbf{A}^{(1)} \odot \mathbf{A}^{(2)}\right)\mathbf{C}^T$ in which $\mathbf{C}^H\mathbf{C} = \mathbf{I}_R$. The orthogonality constrained rank of a tensor $\mathcal{X}$ is equal to the minimal number of orthogonality constrained ($\mathbf{c}_i^H\mathbf{c}_j = \delta_{ij}$) rank-1 terms that yield $\mathcal{X}$ in a linear combination. In [23] uniqueness of CPD with a columnwise orthonormal factor matrix has been demonstrated under milder conditions than an unconstrained CPD. Orthogonality-constrained CPD is very common in statistical signal processing. Typically, the orthonormal columns of $\mathbf{C}$ model signals that are uncorrelated. See [23] for an overview, applications and references. We will now provide a new uniqueness condition for a CPD with a columnwise orthonormal factor matrix.

THEOREM 5.3. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I_1 \times I_2 \times K}$ with matrix representation $\boldsymbol{X}_{(1)} = \left(\boldsymbol{A}^{(1)} \odot \boldsymbol{A}^{(2)}\right)\boldsymbol{C}^T$ in which $\boldsymbol{C}$ is columnwise orthonormal. Let*

$$N_{\max} = \begin{cases} 2 & \text{if} \quad I_2 \geq I_1 \\ 1 & \text{if} \quad I_1 > I_2 \end{cases}, \quad N_{\min} = \begin{cases} 2 & \text{if} \quad I_2 < I_1 \\ 1 & \text{if} \quad I_1 \leq I_2 \end{cases} \tag{5.5}$$

$$I_{\max} = \max(I_1, I_2), \quad I_{\min} = \min(I_1, I_2)\,. \tag{5.6}$$

Construct $\boldsymbol{D} = \boldsymbol{A}^{(N_{\min})*} \odot \boldsymbol{A}^{(N_{\min})} \in \mathbb{C}^{I_{\min}^2 \times R}$. Let S denote a subset of $\{1, \ldots, R\}$ and let $S^c = \{1, \ldots, R\} \setminus S$ denote the complementary set. Stack the columns of $\boldsymbol{D}$ with index in S in $\boldsymbol{D}^{(S)} \in \mathbb{C}^{I_{\min}^2 \times card(S)}$ and stack the columns of $\boldsymbol{D}$ with index in S^c in $\boldsymbol{D}^{(S^c)} \in \mathbb{C}^{I_{\min}^2 \times (R - card(S))}$. Stack the columns of $\boldsymbol{A}^{(N_{\max})}$ in the same order such that $\boldsymbol{A}^{(N_{\max}, S)} \in \mathbb{C}^{I_{\max} \times card(S)}$ and $\boldsymbol{A}^{(N_{\max}, S^c)} \in \mathbb{C}^{I_{\max} \times (R - card(S))}$ are obtained. If

$$C_{R - r_{\boldsymbol{D}} + 2}\left(\boldsymbol{A}^{(N_{\max})*}\right) \odot C_{R - r_{\boldsymbol{D}} + 2}\left(\boldsymbol{A}^{(N_{\max})}\right) \text{ has full column rank,} \tag{5.7a}$$

and if there exists a subset $S \subseteq \{1, \ldots, R\}$ with $0 \leq card(S) \leq r_{\boldsymbol{D}}$ such that

$$\begin{cases} \boldsymbol{D}^{(S)} \text{ has full column rank,} \\ \boldsymbol{A}^{(N_{\max}, S^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{D}^{(S)}}\boldsymbol{D}^{(S^c)}\right) \odot \boldsymbol{A}^{(N_{\max}, S^c)}, \left(\boldsymbol{P}_{\boldsymbol{D}^{(S)}}\boldsymbol{d}_r^{(S^c)}\right) \otimes \boldsymbol{I}_{I_{\max}}\right]\right) = \alpha,\ \forall r \in S^c, \end{cases} \tag{5.7b}$$

where $\alpha = I_{\max} + R - \mathit{card}\,(S) - 1$, *or*

$$r\left(\left[\boldsymbol{A}^{(N_{\min})} \odot \boldsymbol{A}^{(N_{\max})}, \boldsymbol{a}_r^{(N_{\min})} \otimes \boldsymbol{I}_{I_{\max}}\right]\right) = I_{\max} + R - 1,\ \forall r \in \{1, \ldots, R\}, \tag{5.7c}$$

then the orthogonality constrained rank of $\mathcal{X}$ *is* R *and the orthogonality constrained CPD of* $\mathcal{X}$ *is unique.*

Proof. Denote $\mathbf{X}^{(i_1 \cdot \cdot)} = \mathcal{X}(i_1, :, :)$ and construct the fourth-order tensor $\mathcal{Y} \in \mathbb{C}^{I_2 \times I_2 \times I_1 \times I_1}$ with matrix slices $\mathbb{C}^{I_2 \times I_2} \ni \mathbf{Y}^{(\cdot \cdot i_3, i_4)} \triangleq \mathcal{Y}(:, :, i_3, i_4) = \mathbf{X}^{(i_3 \cdot \cdot)} \mathbf{X}^{(i_4 \cdot \cdot) H}$. Since $\mathbf{X}^{(i_1 \cdot \cdot)} = \mathbf{A}^{(2)} D_{i_1}\left(\mathbf{A}^{(1)}\right) \mathbf{C}^T$ and since $\mathbf{C}$ is columnwise orthonormal, the matrix slice $\mathbf{Y}^{(\cdot \cdot i_3, i_4)}$ admits the factorization

$$\mathbf{Y}^{(\cdot \cdot i_3, i_4)} = \mathbf{A}^{(2)} D_{i_3}\left(\mathbf{A}^{(1)}\right) D_{i_4}\left(\mathbf{A}^{(1)*}\right) \mathbf{A}^{(2)H}, \quad i_3, i_4 \in \{1, \ldots, I_1\}.$$

Thus, the fourth-order tensor $\mathcal{Y}$ has the following matrix decomposition

$$\begin{aligned} \mathbf{Y} &= \left[\mathrm{Vec}\left(\mathbf{Y}^{(\cdot \cdot 1,1)}\right), \mathrm{Vec}\left(\mathbf{Y}^{(\cdot \cdot 1,2)}\right), \ldots, \mathrm{Vec}\left(\mathbf{Y}^{(\cdot \cdot I_1, I_1)}\right)\right] \\ &= \left(\mathbf{A}^{(2)*} \odot \mathbf{A}^{(2)}\right)\left(\mathbf{A}^{(1)*} \odot \mathbf{A}^{(1)}\right)^T. \end{aligned} \tag{5.8}$$

If the fourth-order CPD with matrix representation $\mathbf{Y}$ in (5.8) is unique, then the CPD of $\mathcal{X}$ with columnwise orthonormal factor matrix is also unique. Let us interpret (5.8) as a matrix representation of a CPD with factor matrices $\mathbf{A}^{(N_{\max})*}$, $\mathbf{A}^{(N_{\max})}$ and $\mathbf{A}^{(N_{\min})*} \odot \mathbf{A}^{(N_{\min})}$, in which the Khatri-Rao structure of the latter is in first instance ignored.

We first explain that conditions (5.7a) and (5.7b) imply orthogonality constrained CPD uniqueness. Analogous to the proof of Theorem 5.1, condition (5.7b) implies that $k_{\mathbf{D}} \geq 2$. Condition (5.7a) together with Theorem 2.1 now tells us that the rank of $\mathcal{Y}$ is R and the factor matrix $\mathbf{A}^{(N_{\min})*} \odot \mathbf{A}^{(N_{\min})}$ is unique. Since $\mathbf{A}^{(N_{\min})*} \odot \mathbf{A}^{(N_{\min})}$ is unique, $\mathbf{A}^{(N_{\min})}$ is unique. Since $\mathbf{A}^{(N_{\min})}$ is unique and conditions (5.7b) hold, Theorem 4.8 guarantees the uniqueness of the factor matrix $\mathbf{A}^{(N_{\max})}$. The columnwise orthonormal matrix $\mathbf{C}$ follows from $\mathbf{X}_{(1)} = \left(\mathbf{A}^{(1)} \odot \mathbf{A}^{(2)}\right) \mathbf{C}^T$ with $\mathbf{X}_{(1)}$, $\mathbf{A}^{(1)}$ and $\mathbf{A}^{(2)}$ known, via an orthogonal Procrustes problem (e.g. [10]). We can now conclude that the orthogonality constrained CPD of $\mathcal{X}$ is unique and the orthogonality constrained rank of $\mathcal{X}$ is R.

We now explain that conditions (5.7a) and (5.7c) imply orthogonality constrained CPD uniqueness. Analogous to the proof of Theorem 5.1, condition (5.7c) implies that $k_{\mathbf{D}} \geq 2$. Condition (5.7a) together with Theorem 2.1 now tells us that the rank of $\mathcal{Y}$ is R and the factor matrix $\mathbf{A}^{(N_{\min})*} \odot \mathbf{A}^{(N_{\min})}$ is unique. This in turn implies that $\mathbf{A}^{(N_{\min})}$ is unique. Since $\mathbf{A}^{(N_{\min})}$ is unique and conditions (5.7c) hold, Theorem 4.6 guarantees the uniqueness of $\mathbf{A}^{(N_{\max})}$ and $\mathbf{C}$. Thus, the orthogonality constrained CPD of $\mathcal{X}$ is unique and the orthogonality constrained rank of $\mathcal{X}$ is R. □

Theorem 5.3 leads to more relaxed uniqueness conditions than those presented in [23]. For instance, consider the PD of $\mathcal{X} \in \mathbb{C}^{7 \times 4 \times 17}$ with matrix representation $\mathbf{X}_{(1)} = (\mathbf{A} \odot \mathbf{B})\,\mathbf{C}^T$ in which $\mathbf{A} \in \mathbb{C}^{7 \times 17}$, $\mathbf{B} \in \mathbb{C}^{4 \times 17}$ and columnwise orthonormal $\mathbf{C} \in \mathbb{C}^{17 \times 17}$. We randomly generate $\mathbf{A}$, $\mathbf{B}$ and $\mathbf{C}$. For this setting the uniqueness results presented in [23] do not apply. On the other hand, using Lemma 4.1 it can be verified that conditions (5.7a) and (5.7c) in Theorem 5.3 with $\mathrm{card}\,(S) = 9$ are generically satisfied.

6. Numerical Experiments. Let $\mathcal{X} \in \mathbb{C}^{I \times J \times K}$ denote the rank-R tensor of which the CPD is given by (2.1) with $\mathbf{C}$ known. The goal is to estimate $\mathbf{A}$ and $\mathbf{B}$ from the observed tensor $\mathcal{T} = \mathcal{X} + \beta\mathcal{N}$, where $\mathcal{N}$ is an unstructured perturbation tensor and $\beta \in \mathbb{R}$ controls the noise level. The entries of all the involved factor matrices and perturbation tensors are randomly drawn from a Gaussian distribution with zero mean and unit variance.

The following SNR measure will be used: $\text{SNR [dB]} = 10 \log \left(\|\mathbf{X}_{(1)}\|_F^2 / \|\beta \mathbf{N}_{(1)}\|_F^2 \right)$. The distance between a factor matrix, say $\mathbf{A}$, and its estimate $\widehat{\mathbf{A}}$, is measured according to the following criterion: $\mathrm{P}(\mathbf{A}) = \min_\Lambda \|\mathbf{A} - \widehat{\mathbf{A}}\Lambda\|_F / \|\mathbf{A}\|_F$, Λ is a diagonal matrix.

We compare Algorithms 1 and 2 with the ALS method which alternates between the update of the two unknown factor matrices $\mathbf{A}$ and $\mathbf{B}$. See the supplementary material for an efficient implementation of ALS. We consider ALS as a representative of the class of optimization algorithms; for other optimization-based algorithms (e.g. [22]) the conclusions are similar. Let $f_k = \|\mathbf{T}_{(1)} - \widehat{\mathbf{T}}_{(1)}^{(k)}\|_F$, where $\widehat{\mathbf{T}}_{(1)}^{(k)}$ denotes the estimated tensor at iteration k, then we decide that the ALS method has converged when $f_k - f_{k+1} < \epsilon_{\text{ALS}} = 1e-8$ or when the number of iterations exceeds 5000. The ALS method is randomly initialized. Since the best out of ten random initializations is used, the ALS method will be referred to as ALS-10. Algorithms 1 and 2 will be referred to as Alg1 and Alg2, respectively. When Alg1 and Alg2 are followed by at most 500 ALS refinement iterations they will be referred to as Alg1-ALS and Alg2-ALS, respectively. Note that the computational cost of the Alg1 and Alg2 methods is very low; not much more than one ALS iteration. For the choice of the integer $\text{card}(S)$ in Algorithm 2, the value that yields the best fit of $\mathcal{T}$ is retained. More precisely, the integer $\text{card}(S)$ which minimizes $g_{\text{card}(S)} = \|\mathbf{T}_{(1)} - (\widehat{\mathbf{A}}_{\text{card}(S)} \odot \widehat{\mathbf{B}}_{\text{card}(S)})\mathbf{C}^T\|_F$ is retained, where $\widehat{\mathbf{A}}_{\text{card}(S)}$ and $\widehat{\mathbf{B}}_{\text{card}(S)}$ denote the estimates of the factor matrices $\mathbf{A}$ and $\mathbf{B}$ obtained by Algorithm 2 with the given $\text{card}(S)$. Since the data are uniformly distributed we just choose $S = \{1, 2, \ldots, \text{card}(S)\}$ if $\text{card}(S) \geq 1$ (i.e., we pick the first $\text{card}(S)$ columns without trying other combinations) and $S = \emptyset$ if $\text{card}(S) = 0$.

Case 1: $I, K < R$ and $J > R$. The model parameters are $I = 3$, $J = 10$, $K = 4$ and $R = 9$. The existing uniqueness conditions stated in Propositions 3.1 and 3.2 do not apply, indicating a difficult problem. On the other hand, Theorem 4.6 generically guarantees the uniqueness of the decomposition. The mean and standard deviation $\mathrm{P}(\mathbf{A})$ and $\mathrm{P}(\mathbf{B})$ values over 100 trials as function of SNR can be seen in figure 7.1. We notice that the Alg1 and Alg1-ALS methods perform better than the ALS-10 method. Hence, the proposed Alg1 method, possibly followed by few ALS refinement steps, seems to be a good method for computing a CPD with a known factor matrix in cases where only one unknown factor matrix has full column rank.

Case 2: $I, J, K < R$. The model parameters are $I = 4$, $J = 6$, $K = 5$ and $R = 8$. Despite the fact that both Proposition 3.2 and Theorem 4.8 generically guarantee the uniqueness of the decomposition, the problem is difficult since none of the involved factors have full column rank. The mean and standard deviation $\mathrm{P}(\mathbf{A})$ and $\mathrm{P}(\mathbf{B})$ values over 100 trials as a function of SNR can be seen in figure 7.2. On average $\text{card}(S) = 2$, which is the smallest value for which the conditions in Theorem 4.8 are satisfied, yielded the best fit. We notice that in the presence of noise Alg2 performs worse than ALS-10. However, we also observe that Alg2-ALS and ALS-10 perform about the same. Since the Alg2-ALS method is computationally cheaper than the ALS-10 method, Alg2 is an attractive procedure for the initialization of the iterative ALS method. Overall, the Alg2-ALS method seems to be the method of choice for the case where none of the unknown factor matrices have full column rank.

7. Conclusion. Many problems in signal processing can be formulated as tensor decomposition problems with a known factor matrix. We mentioned applications in wireless communication. In the first part of the paper we provided a new uniqueness condition for CPD with a known factor matrix. We showed that by taking the known factor in account more relaxed uniqueness conditions can be obtained compared to unconstrained CPD. We also proposed an inexpensive algebraic method for computing a CPD with a known factor matrix. In the supplementary material an efficient implementation of the ALS method for CPD with a known factor matrix was also reported.

Based on the results obtained in the first part of the paper, we provided in the second part of the paper a new versatile deterministic overall uniqueness condition for the CPD. Since the condition is flexible it can be adapted to other tensor decompositions, as demonstrated in [24, 25]. The supplementary material also contains a partial uniqueness variant of the overall uniqueness result presented in the paper. In tensor-based statistical signal processing CPDs are typically constrained, e.g., orthogonality or Hermitian-symmetry constrained. For that reason we presented new uniqueness conditions for CPD with a partial Hermitian symmetry or columnwise orthonormal factor matrix. Again, such constraints lead in some cases to more relaxed uniqueness conditions than their unconstrained CPD counterparts.

Finally, numerical experiments confirmed the practical use of the proposed Algorithms 1 and 2 for computing a CPD with a known factor matrix. More precisely, in the case where one of the unknown factor matrices had full column rank, Algorithm 1 performed better than the popular ALS method. In the case where none of the unknown factor matrices had full column rank, Algorithm 2 provided at a low computational cost a good initial value for an optimization-based algorithm.

REFERENCES

[1] A. Cichocki, D. Mandic, A.-H. Phan, C. Caifa, G. Zhou, Q. Zhao and L. De Lathauwer, *Tensor decompositions for signal processing applications: From two-way to multiway component analysis*, IEEE Signal Processing Magazine, To appear.

[2] P. Comon and C. Jutten, Editors, *Handbook of blind source separation, independent component analysis and applications*, Academic Press, 2010.

[3] A. L. F. de Almeida, G. Favier and J. C. M. Mota, *Multiuser MIMO systems using space-time-frequency multiple-access PARAFAC tensor modeling*, in: F. Cavalcanti and S. Andersson, editors, Optimizing wireless communication systems, Springer Verlag, 2009.

[4] A. L. F. de Almeida and G. Favier, *Double Khatri-Rao space-time frequency coding using semi-blind PARAFAC based receiver*, IEEE Signal Processing Letters, 10(2013), pp. 471-474.

[5] A. L. F. de Almeida, A. Y. Kibangou, S. Miron and D. C. Araújo, *Joint data and connection topology recovery in collaborative wireless sensor networks*, Proc. ICASSP 2013, 26-31 May, Vancouver, Canada, 2013.

[6] L. De Lathauwer, *A link between the canonical decomposition in multilinear algebra and simultaneous matrix diagonalization*, SIAM J. Matrix Anal. Appl., 28 (2006), pp. 642–666.

[7] L. De Lathauwer, *A short introduction to tensor-based methods for factor analysis and blind source separation*, Proc. ISPA 2011, Sep. 4-6, Dubrovnik, Croatia, 2011.

[8] I. Domanov and L. De Lathauwer, *On the uniqueness of the canonical polyadic decomposition of third-order tensors — Part I: Basic results and uniqueness of one factor matrix*, SIAM J. Matrix Anal. Appl., 34 (2013), pp. 855–875.

[9] I. Domanov and L. De Lathauwer, *On the uniqueness of the canonical polyadic decomposition of third-order tensors — Part II: Overall uniqueness*, SIAM J. Matrix Anal. Appl., 34 (2013), pp. 876–903.

[10] R. A. Horn and C. Johnson, *Matrix analysis*, 2nd ed. Cambridge University Press, 2013.

[11] T. Jiang, N. D. Sidiropoulos and J. M. F. Ten Berge, *Almost-sure identifiability of multidimensional harmonic retrieval*, IEEE Trans. Signal Process., 49(2001), pp. 1849–1859.

[12] T. Jiang and N. D. Sidiropoulos, *Kruskal's permutation lemma and the identification of CANDECOMP/PARAFAC and bilinear models with constant modulus constraints*, IEEE Trans. Signal Process., 52(2004), pp. 2625–2636.
[13] P. M. Kroonenberg, *Applied multiway data analysis*, Wiley, 2008.
[14] T. Kolda and B. W. Bader, *Tensor decompositions and applications*, SIAM Review, 51(2009), 455-500.
[15] J. B. Kruskal, *Three-way arrays: Rank and uniqueness of trilinear decompositions, with applications to arithmetic complexity and statistics*, Linear Algebra and its Applications, 18 (1977), pp. 95–138.
[16] D. Nion and N. D. Sidiropoulos, *Tensor algebra and multidimensional harmonic retrieval in signal processing for MIMO radar*, IEEE Trans. Signal Processing, 58(2010), pp. 5693–5705.
[17] F. Roemer and M. Haardt, *Tensor-based channel estimation and iterative refinements for two-way relaying with multiple antennas and spatial reuse*, IEEE Trans. Signal Processing, 58(2010), pp. 5720-5735.
[18] A. Smilde, R. Bro and P. Geladi, *Multi-way analysis: Applications in the chemical sciences*, Wiley, 2004.
[19] N. D. Sidiropoulos, R. Bro and G. B. Giannakis, *Parallel factor analysis in sensor array processing*, IEEE Trans. Signal Process., 48 (2000), pp. 2377-2388.
[20] N. D. Sidiropoulos, G. B. Giannakis and R.Bro, *Blind PARAFAC receivers for DS-CDMA systems*, IEEE Trans. Signal Process., 48 (2000), pp. 810-823.
[21] N. D. Sidiropoulos and R. S. Budampati, *Khatri-Rao space-time codes*, IEEE Trans. Signal Process., 50 (2002), pp. 2396-2407.
[22] L. Sorber, M. Van Barel and L. De Lathauwer, *Optimization-based algorithms for tensor decompositions: canonical polyadic decomposition, decomposition in rank-(Lr,Lr, 1) terms and a new generalization*, SIAM J. Optim., 23 (2013), pp. 695–720.
[23] M. Sørensen, L. De Lathauwer, P. Comon, S. Icart and L. Deneire, *Canonical polyadic decomposition with a columnwise orthonormal factor matrix*, SIAM J. Matrix Anal. Appl., 33 (2012), pp. 1190–1213.
[24] M. Sørensen and L. De Lathauwer, *Coupled Canonical Polyadic Decompositions and (Coupled) Decompositions in Multilinear rank-$(L_{r,n}, L_{r,n}, 1)$ terms — Part I: Uniqueness*, Internal Report 13-143, ESAT-STADIUS, KU Leuven (Leuven, Belgium), 2013.
[25] M. Sørensen, I. Domanov, D. Nion and L. De Lathauwer, *Coupled canonical polyadic decompositions and (coupled) decompositions in multilinear rank-$(L_{r,n}, L_{r,n}, 1)$ terms — Part II: Algorithms*, Internal Report 13-144, ESAT-STADIUS, KU Leuven (Leuven, Belgium), 2013.
[26] A. Stegeman and N. D. Sidiropoulos, *On Kruskal's uniqueness condition for the Candecomp/Parafac decomposition*, Linear Algebra and its Applications, 420 (2007), pp. 540–552.
[27] A. Stegeman, *On uniqueness of the canonical tensor decomposition with some form of symmetry*, SIAM J. Matrix Anal. Appl., 32(2011), pp. 561–583.
[28] A.-J. Van Der Veen and A. Paulraj, *An analytical constant modulus algorithm*, IEEE Trans. Signal Processing, 44(1996), pp. 1136–1155.



FIG. 7.1. *Mean and standard deviation $P(\boldsymbol{A})$ and $P(\boldsymbol{B})$ for varying SNR, case 1.*



FIG. 7.2. *Mean and standard deviation $P(\boldsymbol{A})$ and $P(\boldsymbol{B})$ for varying SNR, case 2.*

NEW UNIQUENESS CONDITIONS FOR THE CANONICAL POLYADIC DECOMPOSITION OF THIRD-ORDER TENSORS SUPPLEMENTARY MATERIAL

MIKAEL SØRENSEN *† AND LIEVEN DE LATHAUWER *†

S.1. Proof of Lemma 4.3. Given a Vandermonde matrix $\mathbf{A} = [\mathbf{a}_1, \ldots, \mathbf{a}_R] \in \mathbb{C}^{I \times R}$ with distinct generators. Let the column vectors of $\mathbf{U} \in \mathbb{C}^{IJ \times R}$ constitute a basis for range $(\mathbf{A} \odot \mathbf{B})$, where $\mathbf{B} \in \mathbb{C}^{J \times R}$. If the matrix $\underline{\mathbf{A}} \odot \mathbf{B}$ has full column rank, then the matrix

$$\mathbf{G}^{(r)} = [\mathbf{U}, \mathbf{a}_r \otimes \mathbf{I}_J] \in \mathbb{C}^{IJ \times (R+J)} \tag{S.1.1}$$

has rank $R+J-1, \forall r \in \{1, \ldots, R\}$. Generically, $\mathbf{G}^{(r)}$ has rank $R+J-1$ if $(I-1)J+1 \geq R$.

Proof. Consider a Vandermonde matrix $\mathbf{A} \in \mathbb{C}^{I \times R}$ of the form [2, (4.1)] Let us first prove that if $\underline{\mathbf{A}} \odot \mathbf{B}$ has full column rank, then $\mathbf{G}^{(r)}$ has a one-dimensional kernel. Note that all bases for range $(\mathbf{A} \odot \mathbf{B})$ are related via a right multiplication by a nonsingular matrix which does not affect the rank of $\mathbf{G}^{(r)}$. Thus, w.l.o.g. we set $\mathbf{U} = \mathbf{A} \odot \mathbf{B}$ such that $\mathbf{G}^{(r)} \in \mathbb{C}^{IJ \times (R+J)}$ becomes

$$\mathbf{G}^{(r)} = [\mathbf{A} \odot \mathbf{B}, \mathbf{a}_r \otimes \mathbf{I}_J] = \begin{bmatrix} \mathbf{B} & \mathbf{I}_J \\ \mathbf{BZ} & z_r \mathbf{I}_J \\ \vdots & \vdots \\ \mathbf{BZ}^{I-1} & z_r^{I-1} \mathbf{I}_J \end{bmatrix},$$

where $\mathbf{Z} = D_1([z_1, z_2, \ldots, z_R])$. Consider the following nonsingular block-bidiagonal matrix

$$\mathbf{F}^{(r)} = \begin{bmatrix} \mathbf{I}_J & \mathbf{0}_{J,J} & \cdots & \cdots & \mathbf{0}_{J,J} & \mathbf{0}_{J,J} \\ -z_r \mathbf{I}_J & \mathbf{I}_J & \ddots & & \vdots & \vdots \\ \mathbf{0}_{J,J} & -z_r \mathbf{I}_J & \ddots & \ddots & \vdots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \mathbf{0}_{J,J} & \mathbf{0}_{J,J} \\ \mathbf{0}_{J,J} & & \ddots & \ddots & \mathbf{I}_J & \mathbf{0}_{J,J} \\ \mathbf{0}_{J,J} & \cdots & \cdots & \mathbf{0}_{J,J} & -z_r \mathbf{I}_J & \mathbf{I}_J \end{bmatrix} \in \mathbb{C}^{IJ \times IJ},$$

*Group Science, Engineering and Technology, KU Leuven - Kulak, E. Sabbelaan 53, 8500 Kortrijk, Belgium, KU Leuven - E.E. Dept. (ESAT) - STADIUS Center for Dynamical Systems, Signal Processing and Data Analytics, and iMinds Medical IT Department, Kasteelpark Arenberg 10, B-3001 Leuven-Heverlee, Belgium. {Mikael.Sorensen, Lieven.DeLathauwer}@kuleuven-kulak.be.

†Research supported by: (1) Research Council KU Leuven: GOA-MaNet, CoE EF/05/006 Optimization in Engineering (OPTEC), CIF1, STRT1/08/23, (2) F.W.O.: project G.0427.10N, G.0830.14N, G.0881.14N, (3) the Belgian Federal Science Policy Office: IUAP P7 (DYSCO II, Dynamical systems, control and optimization, 2012-2017), (4) EU: ERC advanced grant no. 339804 (BIOTENSORS).

then

$$\mathbf{F}^{(r)}\mathbf{G}^{(r)} = \begin{bmatrix} \mathbf{B} & \mathbf{I}_J \\ \mathbf{B}\left(\mathbf{Z} - z_r\mathbf{I}_R\right) & \mathbf{0}_{J,J} \\ \vdots & \vdots \\ \mathbf{B}\left(\mathbf{Z}^{I-1} - z_r\mathbf{Z}^{I-2}\right) & \mathbf{0}_{J,J} \end{bmatrix}. \tag{S.1.2}$$

Hence, the problem of determining the rank of $\mathbf{G}^{(r)}$ reduces to finding the rank of

$$\begin{aligned} \mathbf{H}^{(r)} &= \begin{bmatrix} \mathbf{B}\left(\mathbf{Z} - z_r\mathbf{I}_R\right) \\ \vdots \\ \mathbf{B}\left(\mathbf{Z}^{I-1} - z_r\mathbf{Z}^{I-2}\right) \end{bmatrix} = \begin{bmatrix} \mathbf{B}\mathbf{I}_R\left(\mathbf{Z} - z_r\mathbf{I}_R\right) \\ \vdots \\ \mathbf{B}\mathbf{Z}^{I-2}\left(\mathbf{Z} - z_r\mathbf{I}_R\right) \end{bmatrix} \\ &= \left(\underline{\mathbf{A}} \odot \mathbf{B}\right)\left(\mathbf{Z} - z_r\mathbf{I}_R\right). \end{aligned} \tag{S.1.3}$$

The rth column vector $\mathbf{h}_r^{(r)}$ of $\mathbf{H}^{(r)}$ is an all-zero vector. Thus the problem of determining the rank of $\mathbf{G}^{(r)}$ further reduces to finding the rank of

$$\widetilde{\mathbf{H}}^{(r)} = \left(\underline{\widetilde{\mathbf{A}}}^{(r)} \odot \widetilde{\mathbf{B}}^{(r)}\right)\widetilde{\mathbf{Z}}^{(r)}, \tag{S.1.4}$$

where

$$\begin{aligned} \widetilde{\mathbf{A}}^{(r)} &= [\mathbf{a}_1, \ldots, \mathbf{a}_{r-1}, \mathbf{a}_{r+1}, \ldots, \mathbf{a}_R] \in \mathbb{C}^{I\times(R-1)}, \\ \widetilde{\mathbf{B}}^{(r)} &= [\mathbf{b}_1, \ldots, \mathbf{b}_{r-1}, \mathbf{b}_{r+1}, \ldots, \mathbf{b}_R] \in \mathbb{C}^{J\times(R-1)}, \\ \widetilde{\mathbf{Z}}^{(r)} &= D_1\left([z_1 - z_r, \ldots, z_{r-1} - z_r, z_{r+1} - z_r, \ldots, z_R - z_r]\right) \in \mathbb{C}^{(R-1)\times(R-1)}. \end{aligned}$$

The deterministic part of the lemma follows from (S.1.4). Indeed, the full column rank assumption on $\underline{\mathbf{A}} \odot \mathbf{B}$ implies that $\underline{\widetilde{\mathbf{A}}}^{(r)} \odot \widetilde{\mathbf{B}}^{(r)}$ has full column rank for every $r \in \{1, \ldots, R\}$. This in turn implies that the matrix in (S.1.3) has rank $R-1$ and consequently $\mathbf{G}^{(r)}$ in (S.1.1) has rank $R+J-1$, $\forall r \in \{1, \ldots, R\}$. We can now conclude that any $\mathbf{U}$ of which the columns form a basis for range$\left(\mathbf{A} \odot \mathbf{B}\right)$ yields a matrix $\mathbf{G}^{(r)}$ of rank $R + J - 1$, $\forall r \in \{1, \ldots, R\}$.

Let us now prove that if $(I-1)J + 1 \geq R$, then generically $\mathbf{G}^{(r)}$ has a one-dimensional kernel. Due to [2, Lemma 4.1] we just need to find one example where the statement made in this lemma holds. Note that any minor of $\mathbf{G}^{(r)}$ is an analytic function of the elements $\mathbf{G}^{(r)}$. Due to [2, Lemma 4.1] we know that when $(I-1)J+1 \geq R$, the matrix $\mathbf{H}^{(r)}$ given by (S.1.3) generically has rank $R-1$. This allows us to conclude that when $(I-1)J+1 \geq R$, the matrix $\mathbf{G}^{(r)}$ in (S.1.1) generically has rank $R + J - 1$, $\forall r \in \{1, \ldots, R\}$. ∎

S.2. Proof of Corollary 4.4. Given $\mathbf{A} = [\mathbf{a}_1, \ldots, \mathbf{a}_R] \in \mathbb{C}^{I\times R}$ and let the column vectors of $\mathbf{U} \in \mathbb{C}^{IJ\times R}$ constitute a basis for range$\left(\mathbf{A} \odot \mathbf{B}\right)$ in which $\mathbf{B} \in \mathbb{C}^{J\times R}$. If $(I-1)J + 1 \geq R$, then the matrix

$$\mathbf{G}^{(r)} = [\mathbf{U}, \mathbf{a}_r \otimes \mathbf{I}_J] \in \mathbb{C}^{IJ\times(R+J)} \tag{S.2.1}$$

generically has rank $R + J - 1$, $\forall r \in \{1, \ldots, R\}$.

Proof. Due to [2, Lemma 4.1] we just need to find one example where the statement made in this corollary holds. By way of example, let $\mathbf{A}$ and $\mathbf{B}$ be Vandermonde matrices and set $\mathbf{U} = \mathbf{A} \odot \mathbf{B}$. Since $(I-1)J+1 \geq R$, the matrix $\mathbf{A} \odot \mathbf{B} \in \mathbb{C}^{IJ \times R}$ generically has full column rank. Due to [2, Lemma 4.3] we know that $\mathbf{G}^{(r)}$ in (S.2.1) generically has rank $R+J-1$, $\forall r \in \{1, \ldots, R\}$. More generally, any $\mathbf{U}$ of which the columns form a basis for range $(\mathbf{A} \odot \mathbf{B})$ yields a matrix $\mathbf{G}^{(r)}$ of rank $R+J-1$, $\forall r \in \{1, \ldots, R\}$. By invoking [2, Lemma 4.1] we can now conclude that when $(I-1)J+1 \geq R$, the matrix $\mathbf{G}^{(r)}$ in (S.2.1) generically has rank $R+J-1$, $\forall r \in \{1, \ldots, R\}$. ☐

S.3. Extension to higher-order tensors with a known factor matrix. Let us briefly explain that the results presented in [2] concerning CPD of a third-order tensor with a known factor can extended to tensors of arbitrary order. Consider the tensor $\mathcal{X} \in \mathbb{C}^{I_1 \times \cdots \times I_P \times J_1 \times \cdots \times J_Q \times K}$ given by

$$\mathcal{X} = \sum_{r=1}^{R} \mathbf{a}_r^{(1)} \circ \cdots \circ \mathbf{a}_r^{(P)} \circ \mathbf{b}_r^{(1)} \circ \cdots \circ \mathbf{b}_r^{(Q)} \circ \mathbf{c}_r$$

with matrix representation

$$\mathbf{X}_{(1)} = \left(\left(\mathbf{A}^{(1)} \odot \cdots \odot \mathbf{A}^{(P)} \right) \odot \left(\mathbf{B}^{(1)} \odot \cdots \odot \mathbf{B}^{(Q)} \right) \right) \mathbf{C}^T = (\mathbf{A} \odot \mathbf{B}) \, \mathbf{C}^T,$$

in which $\mathbf{A}^{(p)} = \left[\mathbf{a}_1^{(p)}, \ldots, \mathbf{a}_R^{(p)} \right] \in \mathbb{C}^{I_p \times R}$, $\mathbf{B}^{(q)} = \left[\mathbf{b}_1^{(q)}, \ldots, \mathbf{b}_R^{(q)} \right] \in \mathbb{C}^{J_q \times R}$, $\mathbf{C} = [\mathbf{c}_1, \ldots, \mathbf{c}_R] \in \mathbb{C}^{K \times R}$, $\mathbf{A} = \mathbf{A}^{(1)} \odot \cdots \odot \mathbf{A}^{(P)}$ and $\mathbf{B} = \mathbf{B}^{(1)} \odot \cdots \odot \mathbf{B}^{(Q)}$. Assume that $\mathbf{C}$ is known.

S.3.1. Uniqueness condition. By ignoring the Khatri-Rao structure within $\mathbf{A}$ and $\mathbf{B}$ we may treat $\mathcal{X}$ as a third-order tensor with dimensions $I = \prod_{p=1}^{P} I_p$, $J = \prod_{q=1}^{Q} J_q$ and K. This in turn means that conditions (4.25a)/(4.25b) in [2] also yield a deterministic condition for the CPD of $\mathcal{X}$ with $\mathbf{C}$ known. Furthermore, it can be shown that Lemma 4.3 and Proposition 4.7 in [2] can be extended to tensors of arbitrary order with a known factor. Consequently, condition (4.26) in [2] also yields a generic condition for the CPD of $\mathcal{X}$ of arbitrary order with a known factor $\mathbf{C}$.

S.3.2. Algorithm. We now explain how to extend Algorithm 2 in [2] to tensors of arbitrary order. As in Theorem 4.8 in [2] S denotes a subset of $\{1, \ldots, R\}$ and $S^c = \{1, \ldots, R\} \setminus S$ denotes the complementary set. Stack the columns of $\mathbf{C}$ with index in S in $\mathbf{C}^{(S)} \in \mathbb{C}^{K \times \mathrm{card}(S)}$ and stack the columns of $\mathbf{C}$ with index in S^c in $\mathbf{C}^{(S^c)} \in \mathbb{C}^{K \times (R - \mathrm{card}(S))}$. Stack the columns of $\mathbf{A}$ (resp. $\mathbf{B}$) in the same order such that $\mathbf{A}^{(S)} \in \mathbb{C}^{I \times \mathrm{card}(S)}$ (resp. $\mathbf{B}^{(S)} \in \mathbb{C}^{J \times \mathrm{card}(S)}$) and $\mathbf{A}^{(S^c)} \in \mathbb{C}^{I \times (R - \mathrm{card}(S))}$ (resp. $\mathbf{B}^{(S^c)} \in \mathbb{C}^{J \times (R - \mathrm{card}(S))}$) are obtained. As in the third-order case, we first compute $\mathbf{Y}_{(1)} = \mathbf{X}_{(1)} \mathbf{P}_{\mathbf{C}^{(S)}}^T$. Denote $\mathbf{D}^{(S^c)} = \mathbf{P}_{\mathbf{C}^{(S)}} \mathbf{C}^{(S^c)}$. Let the columns of $\mathbf{U}$ constitute a basis for the range of $\mathbf{Y}_{(3)} = \left(\mathbf{D}^{(S^c)} \odot \mathbf{A}^{(S^c)} \right) \mathbf{B}^{(S^c)T}$. If the conditions stated in [2, Theorem 4.8] are satisfied, then there exists a nonsingular matrix $\mathbf{M}$ such that

$$\mathbf{UM} = \mathbf{D}^{(S^c)} \odot \mathbf{A}^{(S^c)} \Leftrightarrow \mathbf{G}^{(r,S^c)} \begin{bmatrix} \mathbf{m}_r \\ \mathbf{a}_r^{(1,S^c)} \otimes \cdots \otimes \mathbf{a}_r^{(P,S^c)} \end{bmatrix} = \mathbf{0}, \quad r \in S^c, \quad \text{(S.3.1)}$$

where

$$\mathbf{G}^{(r,S^c)} = \left[\mathbf{U}, -\mathbf{d}_r^{(S^c)} \otimes \mathbf{I}_I \right]. \quad \text{(S.3.2)}$$

We also know that if the conditions stated in [2, Theorem 4.8] are satisfied, then the matrices $\mathbf{G}^{(r,S^c)}$ given by (S.3.2) have a one-dimensional kernel. This means that by solving the set of linear systems $\mathbf{G}^{(r,S^c)}\mathbf{z}_r = \mathbf{0}$, $r \in S^c$, we can obtain $\mathbf{A}^{(S^c)}$ from the solutions $\{\mathbf{z}_r\}$ via best rank-1 approximations. The remaining steps are analogous to the third-order case. We summarize the procedure for computing the CPD of a higher-order tensor with a known factor matrix in Algorithm 1.

Algorithm 1 Outline of the procedure for computing a CPD with a known factor matrix $\mathbf{C}$ based on an extension of Theorem 4.8 in [2] to tensors of arbitrary order.

Input: $\mathbf{X}_{(1)} = \left(\left(\mathbf{A}^{(1)} \odot \cdots \odot \mathbf{A}^{(P)}\right) \odot \left(\mathbf{B}^{(1)} \odot \cdots \odot \mathbf{B}^{(Q)}\right)\right)\mathbf{C}^T$ and $\mathbf{C}$.

1. Determine $r_{\mathbf{C}} = r(\mathbf{C})$.
2. Choose sets $S \subseteq \{1,\ldots,R\}$ and $S^c = \{1,\ldots,R\}\backslash S$ subject to $0 \leq \operatorname{card}(S) \leq r_{\mathbf{C}}$.
3. Build $\mathbf{C}^{(S)}$ and $\mathbf{C}^{(S^c)}$.
4. Find $\mathbf{F}$ whose column vectors constitute an orthonormal basis for range $\left(\mathbf{C}^{(S)}\right)$.
5. Compute $\mathbf{P}_{\mathbf{C}^{(S)}} = \mathbf{I}_K - \mathbf{F}\mathbf{F}^H$ and $\mathbf{D}^{(S^c)} = \mathbf{P}_{\mathbf{C}^{(S)}}\mathbf{C}^{(S^c)}$.
6. Compute $\mathbf{Y}_{(1)} = \mathbf{X}_{(1)}\mathbf{P}^T_{\mathbf{C}^{(S)}}$.
7. Find $\mathbf{U}$ whose column vectors constitute an orthonormal basis for range $\left(\mathbf{Y}_{(3)}\right)$.
8. Set $I = \prod_{p=1}^{P} I_p$.
9. Solve set of linear equations $\left[\mathbf{U}, -\mathbf{d}_r^{(S^c)} \otimes \mathbf{I}_I\right]\mathbf{z}_r = \mathbf{0}_{KI}, \quad r \in S^c$.
10. Set $\mathbf{a}_r^{(S^c)} = \mathbf{z}_r(R+1:R+I), \quad r \in S^c$.
11. Solve rank-1 approximations
$$\min_{\mathbf{a}_r^{(1,S^c)},\ldots,\mathbf{a}_r^{(P,S^c)}} \left\|\mathbf{a}_r^{(S^c)} - \mathbf{a}_r^{(1,S^c)} \otimes \cdots \otimes \mathbf{a}_r^{(P,S^c)}\right\|_F^2, \quad r \in S^c.$$
12. Compute $\mathbf{G}^{(S^c)} = \left(\mathbf{A}^{(1,S^c)H}\mathbf{A}^{(1,S^c)}\right) * \cdots * \left(\mathbf{A}^{(P,S^c)H}\mathbf{A}^{(P,S^c)}\right)$.
13. Compute $\mathbf{H}^{(S^c)} = \mathbf{A}^{(1,S^c)} \odot \cdots \odot \mathbf{A}^{(P,S^c)}$.
14. Compute $\mathbf{B}^{(S^c)T} = \left(\left(\mathbf{C}^{(S^c)H}\mathbf{D}^{(S^c)}\right) * \mathbf{G}^{(S^c)}\right)^{-1}\left(\mathbf{D}^{(S^c)} \odot \mathbf{H}^{(S^c)}\right)^H \mathbf{Y}_{(3)}$.
15. Solve rank-1 approximations
$$\min_{\mathbf{b}_r^{(1,S^c)},\ldots,\mathbf{b}_r^{(Q,S^c)}} \left\|\mathbf{b}_r^{(S^c)} - \mathbf{b}_r^{(1,S^c)} \otimes \cdots \otimes \mathbf{b}_r^{(Q,S^c)}\right\|_F^2, \quad r \in S^c.$$
16. Compute $\mathbf{Q}_{(1)} = \mathbf{X}_{(1)} - \left(\mathbf{H}^{(S^c)} \odot \mathbf{B}^{(1,S^c)} \odot \cdots \odot \mathbf{B}^{(Q,S^c)}\right)\mathbf{C}^{(S^c)T}$.
17. Compute $\mathbf{H} = \mathbf{Q}_{(1)}\left(\mathbf{C}^{(S)T}\right)^{\dagger}$.
18. Solve rank-1 approximations
$$\min_{\substack{\mathbf{a}_r^{(1,S)},\ldots,\mathbf{a}_r^{(P,S)} \\ \mathbf{b}_r^{(1,S)},\ldots,\mathbf{b}_r^{(Q,S)}}} \left\|\mathbf{h}_r - \mathbf{a}_r^{(1,S)} \otimes \cdots \otimes \mathbf{a}_r^{(P,S)} \otimes \mathbf{b}_r^{(1,S)} \otimes \cdots \otimes \mathbf{b}_r^{(Q,S)}\right\|_F^2, \; r \in S.$$

Output: $\{\mathbf{A}^{(p)}\}_{p=1}^P$ and $\{\mathbf{B}^{(q)}\}_{q=1}^Q$.

S.4. Generic rank of $\mathbf{D} = [\mathbf{C}^T, \mathbf{C}^H]^T$. Consider a generic matrix $\mathbf{C} \in \mathbb{C}^{K\times R}$. The matrix
$$\mathbf{D} = \begin{bmatrix} \mathbf{C} \\ \mathbf{C}^* \end{bmatrix} \in \mathbb{C}^{2K\times R}$$
generically has rank $\min(2K, R)$.

Proof. The result is straightforward, simply notice that

$$\begin{bmatrix} \mathbf{C} \\ \mathbf{C}^* \end{bmatrix} = \begin{bmatrix} \mathbf{I}_K & \sqrt{-1}\cdot\mathbf{I}_K \\ \mathbf{I}_K & -\sqrt{-1}\cdot\mathbf{I}_K \end{bmatrix} \begin{bmatrix} \mathrm{Re}\{\mathbf{C}\} \\ \mathrm{Im}\{\mathbf{C}\} \end{bmatrix} = \mathbf{T} \begin{bmatrix} \mathrm{Re}\{\mathbf{C}\} \\ \mathrm{Im}\{\mathbf{C}\} \end{bmatrix}.$$

Due to [2, Lemma 4.1] we just need to find one example where the statement made in this proposition holds. By way of example, if $2K \geq R$, we let $\left[\mathrm{Re}\{\mathbf{C}\}^T, \mathrm{Im}\{\mathbf{C}\}^T\right]^T = \left[\mathbf{I}_R^T, \mathbf{0}_{2K-R,R}^T\right]^T$ and if $2K < R$, we let $\left[\mathrm{Re}\{\mathbf{C}\}^T, \mathrm{Im}\{\mathbf{C}\}^T\right]^T = \left[\mathbf{I}_{2K}^T, \mathbf{0}_{2K,R-2K}^T\right]^T$. Since $\mathbf{T} \in \mathbb{C}^{2K\times 2K}$ is nonsingular we know that for this particular case the rank of $\left[\mathbf{C}^T, \mathbf{C}^H\right]^T$ is equal to $\min(2K, R)$. By invoking [2, Lemma 4.1] we conclude that $\left[\mathbf{C}^T, \mathbf{C}^H\right]^T$ generically has rank $\min(2K, R)$. □

S.5. A new partial uniqueness condition for the CPD. Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$:

$$\mathcal{X} = \sum_{r=1}^{R} \mathbf{a}_r \circ \mathbf{b}_r \circ \mathbf{c}_r \,. \tag{S.5.1}$$

In [1] a partial uniqueness condition for the CPD of $\mathcal{X}$ was obtained (Theorem 3.1 in [1], given below as Lemma S.5.1). Partial uniqueness means that when the overall CPD is not unique, some submatrices of the factor matrices may still be unique. Theorem S.5.2 presents a new partial uniqueness condition obtained from [2, Theorem 4.8]. The derivation will make use of the following lemma.

LEMMA S.5.1. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in* (S.5.1). *Assume that $\boldsymbol{A}$ is unique and that there exists a permutation matrix $\Pi \in \mathbb{C}^{R\times R}$ such that the partitioning*

$$\boldsymbol{A}\Pi = \left[\boldsymbol{A}^{(1)}, \ldots, \boldsymbol{A}^{(Q)}\right], \quad \boldsymbol{A}^{(q)} \in \mathbb{C}^{I\times R_q}, \tag{S.5.2}$$

$$\boldsymbol{B}\Pi = \left[\boldsymbol{B}^{(1)}, \ldots, \boldsymbol{B}^{(Q)}\right], \quad \boldsymbol{B}^{(q)} \in \mathbb{C}^{J\times R_q}, \tag{S.5.3}$$

$$\boldsymbol{C}\Pi = \left[\boldsymbol{C}^{(1)}, \ldots, \boldsymbol{C}^{(Q)}\right], \quad \boldsymbol{C}^{(q)} \in \mathbb{C}^{K\times R_q}, \tag{S.5.4}$$

where $R = \sum_{q=1}^{Q} R_q$ has the following property

$$\mathit{range}\left(\boldsymbol{A}^{(q)}\right) \bigcap \left(\bigcup_{\substack{p=1 \\ p\neq q}}^{Q} \mathit{range}\left(\boldsymbol{A}^{(p)}\right) \right) = \{\boldsymbol{0}_I\}. \tag{S.5.5}$$

Then the pair $\left(\boldsymbol{B}^{(q)}, \boldsymbol{C}^{(q)}\right)$ is unique [1].

Note that if $R_q = 1$, then due to the rank-1 structure of $\mathbf{X}_{(1)}^{(q)} = \left(\mathbf{a}^{(q)} \otimes \mathbf{b}^{(q)}\right)\mathbf{c}^{(q)T}$ the pair $\left(\mathbf{b}^{(q)}, \mathbf{c}^{(q)}\right)$ is obviously unique when $\mathbf{a}^{(q)}$ is given. Consequently, in the following Theorem S.5.2 we will only consider the case where $R_q \geq 2$.

THEOREM S.5.2. *Consider the PD of $\mathcal{X} \in \mathbb{C}^{I\times J\times K}$ in* (S.5.1). *Let $\Pi \in \mathbb{C}^{R\times R}$ be a permutation matrix such that $\boldsymbol{A}$ has a partitioning* (S.5.2) *with property* (S.5.5). *The corresponding partitionings of $\boldsymbol{B}$ and $\boldsymbol{C}$ are denoted by* (S.5.3) *and* (S.5.4), *respectively.*

Let $S_q \subseteq \{1,\ldots,R_q\}$ denote a subset of $\{1,\ldots,R_q\}$ and let $S_q^c = \{1,\ldots,R_q\} \setminus S_q$ denote the complementary set. Stack the columns of $\boldsymbol{C}^{(q)}$ with index in S_q in $\boldsymbol{C}^{(q,S_q)} \in \mathbb{C}^{K \times card(S_q)}$ and stack the columns of $\boldsymbol{C}^{(q)}$ with index in S_q^c in $\boldsymbol{C}^{(q,S_q^c)} \in \mathbb{C}^{K \times (R_q - card(S_q))}$. Stack the columns of $\boldsymbol{A}^{(q)}$ (resp. $\boldsymbol{B}^{(q)}$) in the same order such that $\boldsymbol{A}^{(q,S_q)} \in \mathbb{C}^{I \times card(S_q)}$ (resp. $\boldsymbol{B}^{(q,S_q)} \in \mathbb{C}^{J \times card(S_q)}$) and $\boldsymbol{A}^{(q,S^c)} \in \mathbb{C}^{I \times (R_q - card(S_q))}$ (resp. $\boldsymbol{B}^{(q,S_q^c)} \in \mathbb{C}^{J \times (R_q - card(S_q))}$) are obtained. If

$$\begin{cases} k_{\boldsymbol{A}} \geq 1, \\ C_{R-r_{\boldsymbol{A}}+2}\left(\boldsymbol{B}\right) \odot C_{R-r_{\boldsymbol{A}}+2}\left(\boldsymbol{C}\right) \text{ has full column rank,} \end{cases} \tag{S.5.6}$$

and if there exists a subset $S_q \subseteq \{1,\ldots,R_q\}$ with $0 \leq card(S_q) \leq r_{\boldsymbol{A}^{(q)}}$ such that $\boldsymbol{A}^{(q,S_q)}$ has full column rank and

$$\begin{cases} \boldsymbol{C}^{(q,S_q^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{A}^{(q,S_q)}} \boldsymbol{A}^{(q,S_q^c)}\right) \odot \boldsymbol{B}^{(q,S_q^c)}, \left(\boldsymbol{P}_{\boldsymbol{A}^{(q,S_q)}} \boldsymbol{a}_{r_q}^{(q,S_q^c)}\right) \otimes \boldsymbol{I}_J\right]\right) = \alpha_q, \, \forall r \in S_q^c, \end{cases} \tag{S.5.7a}$$

where $\alpha_q = J + R_q - card(S_q) - 1$, or

$$\begin{cases} \boldsymbol{B}^{(q,S_q^c)} \text{ has full column rank,} \\ r\left(\left[\left(\boldsymbol{P}_{\boldsymbol{A}^{(q,S_q)}} \boldsymbol{A}^{(q,S_q^c)}\right) \odot \boldsymbol{C}^{(q,S_q^c)}, \left(\boldsymbol{P}_{\boldsymbol{A}^{(q,S_q)}} \boldsymbol{a}_{r_q}^{(q,S_q^c)}\right) \otimes \boldsymbol{I}_K\right]\right) = \beta_q, \, \forall r \in S_q^c, \end{cases} \tag{S.5.7b}$$

where $\beta_q = K + R_q - card(S_q) - 1$, then the rank of $\mathcal{X}$ is R, $\boldsymbol{A}$ is unique and the pair $\left(\boldsymbol{B}^{(q)}, \boldsymbol{C}^{(q)}\right)$ is unique.

Proof. Condition (S.5.6) implies that the rank of $\mathcal{X}$ is R and the factor matrix $\mathbf{A}$ is unique [2, Theorem 2.1]. Lemma S.5.1 tells us that the study of the overall uniqueness of $\mathcal{X}$ decouples into the study of the CPDs with matrix representations

$$\mathbf{X}_{(1)}^{(q)} = \left(\mathbf{A}^{(q)} \odot \mathbf{B}^{(q)}\right) \mathbf{C}^{(q)T}, \quad q \in \{1,\ldots,Q\},$$

where $\mathbf{A}^{(q)} \in \mathbb{C}^{I \times R_q}$, $\mathbf{B}^{(q)} \in \mathbb{C}^{J \times R_q}$ and $\mathbf{C}^{(q)} \in \mathbb{C}^{K \times R_q}$ are given by (S.5.2), (S.5.3) and (S.5.4), respectively. Fix $\mathbf{A}^{(q)}$, then [2, Theorem 4.8] together with the conditions (S.5.7a)/(S.5.7b) now guarantee the uniqueness of the pair $\left(\mathbf{B}^{(q)}, \mathbf{C}^{(q)}\right)$. ∎

S.6. ALS method for CPD with a known factor matrix. Consider the PD of $\mathcal{X} \in \mathbb{C}^{I \times J \times K}$ with matrix representation $\mathbf{X}_{(2)} = (\mathbf{B} \odot \mathbf{C})\,\mathbf{A}^T$ in which $\mathbf{C}$ is known. The following identity

$$(\mathbf{B} \odot \mathbf{C})^\dagger = \left(\left(\mathbf{B}^H\mathbf{B}\right) * \left(\mathbf{C}^H\mathbf{C}\right)\right)^{-1} (\mathbf{B} \odot \mathbf{C})^H$$

is well-known. Hence, the conditional least squares update of $\mathbf{A}$ while $\mathbf{B}$ is fixed is

$$\mathbf{A}^T = (\mathbf{B} \odot \mathbf{C})^\dagger \mathbf{X}_{(2)} = \left(\left(\mathbf{B}^H\mathbf{B}\right) * \left(\mathbf{C}^H\mathbf{C}\right)\right)^{-1} (\mathbf{B} \odot \mathbf{C})^H \mathbf{X}_{(2)}. \tag{S.6.1}$$

Using[1]

$$\left(\mathbf{B}\odot\mathbf{C}\right)^{H}\mathbf{X}_{(2)}(:,i)=\mathrm{Vecd}\left(\mathbf{B}^{H}\mathbf{X}^{(i\cdot\cdot)}\mathbf{C}^{*}\right)=\begin{bmatrix}\mathbf{b}_1^{H}\mathbf{X}^{(i\cdot\cdot)}\mathbf{c}_1^{*}\\ \vdots\\ \mathbf{b}_R^{H}\mathbf{X}^{(i\cdot\cdot)}\mathbf{c}_R^{*}\end{bmatrix}=\begin{bmatrix}\mathbf{b}_1^{H}\mathbf{y}_i^{(1)}\\ \vdots\\ \mathbf{b}_R^{H}\mathbf{y}_i^{(R)}\end{bmatrix},$$

where $\mathbf{y}_i^{(r)}=\mathbf{X}^{(i\cdot\cdot)}\mathbf{c}_r^{*}$, relation (S.6.1) can be expressed as follows:

$$\mathbf{A}^{T}=\left(\mathbf{B}\odot\mathbf{C}\right)^{\dagger}\mathbf{X}_{(2)}=\left(\left(\mathbf{B}^{H}\mathbf{B}\right)*\left(\mathbf{C}^{H}\mathbf{C}\right)\right)^{-1}\begin{bmatrix}\mathbf{b}_1^{H}\mathbf{Y}^{(1)}\\ \vdots\\ \mathbf{b}_R^{H}\mathbf{Y}^{(R)}\end{bmatrix},$$

where $\mathbf{Y}^{(r)}=\left[\mathbf{y}_1^{(r)},\ldots,\mathbf{y}_I^{(r)}\right]$. Similarly, by making use of

$$\left(\mathbf{C}\odot\mathbf{A}\right)^{\dagger}=\left(\left(\mathbf{C}^{H}\mathbf{C}\right)*\left(\mathbf{A}^{H}\mathbf{A}\right)\right)^{-1}\left(\mathbf{C}\odot\mathbf{A}\right)^{H},$$

$$\left(\mathbf{C}\odot\mathbf{A}\right)^{H}\mathbf{X}_{(3)}(:,j)=\mathrm{Vecd}\left(\mathbf{A}^{H}\mathbf{X}^{(\cdot j\cdot)}\mathbf{C}^{*}\right)=\begin{bmatrix}\mathbf{a}_1^{H}\mathbf{X}^{(\cdot j\cdot)}\mathbf{c}_1^{*}\\ \vdots\\ \mathbf{a}_R^{H}\mathbf{X}^{(\cdot j\cdot)}\mathbf{c}_R^{*}\end{bmatrix}=\begin{bmatrix}\mathbf{a}_1^{H}\mathbf{z}_j^{(1)}\\ \vdots\\ \mathbf{a}_R^{H}\mathbf{z}_j^{(R)}\end{bmatrix},$$

where $\mathbf{z}_j^{(r)}=\mathbf{X}^{(\cdot j\cdot)}\mathbf{c}_r^{*}$, the conditional least squares update of $\mathbf{B}$ can be expressed as

$$\mathbf{B}^{T}=\left(\mathbf{C}\odot\mathbf{A}\right)^{\dagger}\mathbf{X}_{(3)}=\left(\left(\mathbf{C}^{H}\mathbf{C}\right)*\left(\mathbf{A}^{H}\mathbf{A}\right)\right)^{-1}\begin{bmatrix}\mathbf{a}_1^{H}\mathbf{Z}^{(1)}\\ \vdots\\ \mathbf{a}_R^{H}\mathbf{Z}^{(R)}\end{bmatrix},$$

where $\mathbf{Z}^{(r)}=\left[\mathbf{z}_1^{(r)},\ldots,\mathbf{z}_J^{(r)}\right]$. We summarize the ALS method for computing a CPD with a known factor matrix in Algorithm 2. The normalization in step 3 fixes the scaling ambiguity. The extension to tensors of order higher than three is straightforward.

REFERENCES

[1] X. Guo, S. Miron, D. Brie and A. Stegeman, *Uni-mode and partial uniqueness conditions for CANDECOMP/PARAFAC of three-way arrays with linearly dependent loadings*, SIAM J. Matrix Anal. Appl., 33 (2012), pp. 111–129.

[2] M. Sørensen and L. De Lathauwer, *New uniqueness conditions for the canonical polyadic decomposition of third-order tensors*, Internal Report 13-05, ESAT-STADIUS, KU Leuven, Belgium.

[1] Let $\mathbf{A}\in\mathbb{C}^{I\times I}$, then $\mathrm{Vecd}\left(\mathbf{A}\right)\in\mathbb{C}^{I}$ denotes the column vector defined by $\left(\mathrm{Vecd}\left(\mathbf{A}\right)\right)_i=\left(\mathbf{A}\right)_{ii}$.

Algorithm 2 ALS procedure for CPD with known factor matrix $\mathbf{C}$.

Input: $\mathbf{X}_{(1)} = (\mathbf{A} \odot \mathbf{B})\,\mathbf{C}^T$ and $\mathbf{C}$.

Initialize $\mathbf{A}$ and $\mathbf{B}$.

Compute $\mathbf{D} = \mathbf{C}^H\mathbf{C}$.

Compute $\mathbf{Y}^{(r)} = \left[\mathbf{y}_1^{(r)}, \ldots, \mathbf{y}_I^{(r)}\right]$ with $\mathbf{y}_i^{(r)} = \mathbf{X}^{(i\cdot\cdot)}\mathbf{c}_r^*, \quad r \in \{1, \ldots, R\}$.

Compute $\mathbf{Z}^{(r)} = \left[\mathbf{z}_1^{(r)}, \ldots, \mathbf{z}_J^{(r)}\right]$ with $\mathbf{z}_j^{(r)} = \mathbf{X}^{(\cdot j\cdot)}\mathbf{c}_r^*, \quad r \in \{1, \ldots, R\}$.

Repeat until convergence:

1. Compute $\mathbf{A}^T = \left(\left(\mathbf{B}^H\mathbf{B}\right) * \mathbf{D}\right)^{-1} \begin{bmatrix} \mathbf{b}_1^H\mathbf{Y}^{(1)} \\ \vdots \\ \mathbf{b}_R^H\mathbf{Y}^{(R)} \end{bmatrix}$.
2. Compute $\mathbf{B}^T = \left(\mathbf{D} * \left(\mathbf{A}^H\mathbf{A}\right)\right)^{-1} \begin{bmatrix} \mathbf{a}_1^H\mathbf{Z}^{(1)} \\ \vdots \\ \mathbf{a}_R^H\mathbf{Z}^{(R)} \end{bmatrix}$.
3. Update $\mathbf{b}_r \leftarrow \frac{\mathbf{b}_r}{\|\mathbf{b}_r\|_F}, \quad r \in \{1, \ldots, R\}$.

Output: $\mathbf{A}$ and $\mathbf{B}$.